

# Origin of powdery mildew resistance factors in RE714, a wheat breeding line obtained from two interspecific crosses

H. Muranty, M.T. Pavoine, G. Doussinault, Dominique Barloy

► **To cite this version:**

H. Muranty, M.T. Pavoine, G. Doussinault, Dominique Barloy. Origin of powdery mildew resistance factors in RE714, a wheat breeding line obtained from two interspecific crosses. *Plant Breeding*, Wiley, 2010, 129 (5), pp.465-471. 10.1111/j.1439-0523.2009.01740.x . hal-00729590

**HAL Id: hal-00729590**

**<https://hal-agrocampus-ouest.archives-ouvertes.fr/hal-00729590>**

Submitted on 31 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Origin of powdery mildew genetic resistance factors in RE714, a wheat breeding line obtained from two interspecific crosses

H. Muranty<sup>1,2</sup>, M.-T. Pavoine<sup>1</sup>, G. Doussinault<sup>1</sup>, D. Barloy<sup>1</sup>

<sup>1</sup>INRA Agrocampus Rennes UMR118 Amélioration des Plantes et Biotechnologies  
Végétales, BP 35327, 35653 LE RHEU CEDEX, FRANCE

<sup>2</sup>Present address : INRA UR 0588 Amélioration Génétique et Physiologie Forestières,  
2163 avenue de la Pomme de Pin - CS40001 - Ardon, 45075 ORLEANS CEDEX 2,  
FRANCE. [Helene.Muranty@orleans.inra.fr](mailto:Helene.Muranty@orleans.inra.fr), phone: 33 2 38 41 78 47, fax : 33 2 38 41 78 79

With 2 figures and 3 tables

Received July 10, 2009/ Accepted October 24, 2009

Communicated by R.A. McIntosh

### Abstract

RE714 is a winter wheat breeding line obtained from two interspecific crosses [(*Aegilops tauschii* no. 33/*Triticum dicoccum* no. 119) and VPM = (*Aegilops ventricosa* no.10/*T. persicum*)]//Marne]. RE714 expresses powdery mildew resistance at the seedling, vernalized and adult stages. The origin of the main genetic factors involved in RE714 powdery mildew resistance (*Pm4b*, *QPm.inra-5D*, *QPm.inra-6A2*, *QPm.inra-7A3a* and *QPm.inra-7A3b*) was studied by comparing alleles of RE714 and those of seven of its parental lines with 36 microsatellite markers previously detected as linked to these factors. VPM, and among its parental lines, the *T. persicum* accession, were identified as the donor of *Pm4b* to RE714. *Aegilops tauschii* n°33 and *T. dicoccum* n°119 were identified as the donors of *QPm.inra-5D* and *QPm.inra-6A2* to RE714, respectively. Beauchamp, a line used at the end of the breeding scheme that created RE714, might be the donor of the favourable alleles of RE714 at quantitative trait loci on linkage group 7A3. These results are discussed in the context of introgression of valuable genetic factors from related species of the primary gene pool into wheat.

**Key words:** wheat – introgression – powdery mildew – resistance genes

Domestication and modern intensive plant breeding are generally considered to be practices that lead to reduced genetic diversity in crop cultivars. For example, Roussel et al. (2004) showed a decrease in allelic richness of about 25% between landraces cultivated between 1800 and 1840 and varieties cultivated after 1840 for common wheat (*Triticum aestivum* L) in France. Wild relatives and related species of crop plants represent the best hope for continuous crop genetic improvement because they harbour desirable agronomic traits such as biotic and abiotic stress resistances and special quality traits (Xie and Nevo 2008). Awareness of this fact led to the building of several conservation programmes of plant genetic resources, either *ex situ* or *in situ*. Although conservation of biodiversity is still needed, the next challenge is to develop innovative breeding strategies to incorporate favourable alleles, genes or gene complexes from genetic resources into cultivated genotypes (Feuillet et al. 2008) without introducing too much associated wild genetic material with unacceptable wild traits. Associated unfavourable genetic material is often referred to as linkage drag. Genetic recombination during meiosis followed by selection can be used to reduce linkage drag. For about 10 years, breeders have introduced molecular markers in the selection step to tag the favourable allele(s) and to estimate the length of the associated introduced segment(s) and reduce it in marker-assisted backcross or genotype building schemes (Hospital 2005).

The most cultivated wheat species today are bread wheat (*Triticum aestivum*) and pasta wheat (*T. durum*), which are hexaploid (AABBDD) and tetraploid (AABB), respectively. The wild relatives of wheat can be classified on the basis of their genomic constitutions into primary, secondary and tertiary gene pools (Jiang et al. 1994). The primary gene pool includes landraces, early domesticates and wild relatives that hybridize directly with the cultivated types. Their genomes are homologous to the AA (*T. monococcum*, including var. *beoticum* and var. *urartu*) or the DD (*Aegilops tauschii*) or the AABB genomes (*T. turgidum*, *T. dicoccum* and *T. dicoccoides*) of cultivated wheats. The secondary gene pool contains polyploid species that share at least one homologous genome with the cultivated types. The tertiary gene pool contains species that do not share a homologous genome with the cultivated types. Gene transfer from these species can only be achieved by sophisticated strategies generally referred to as "chromosome engineering" (Qi et al. 2007).

The winter wheat breeding line RE714 was developed from an interspecific cross (*Aegilops tauschii* no. 33/*Triticum dicoccum* no. 119) which was crossed to several lines including lines derived from VPM/Moisson (VPM = *Aegilops ventricosa* no.10/*T. persicum*//3\* Marne) in a breeding program for disease resistance at INRA, Rennes (France) (Trottet and Dosba 1983; Robe and Doussinault 1995); (Fig. 1). These crosses involved species from the primary gene

pool of cultivated bread wheat, namely *Ae. tauschii*, with the D genome, and *T. dicoccum* and *T. persicum* (= *T. carthlicum*), with the A and B genomes, and a species from the secondary gene pool, namely *Ae. ventricosa*, with the D and N<sup>v</sup> genomes. G. Doussinault sent seeds of RE714 to several breeders in France and Europe, but few reported if they used it: we only know that it was used in Switzerland (D. Fossati pers. comm.). RE714 expresses powdery mildew resistance at the seedling, vernalized and adult stage and the genetic basis of its resistance has been studied in detail (Robe and Doussinault 1995, Chantret et al. 1999, 2000, 2001, Mingeot et al. 2002, Muranty et al. 2008, 2009). Three main genetic factors control RE714 powdery mildew resistance at the seedling stage. These are *Pm4b*, located on chromosome 2A, *QPms.inra-5D* located on chromosome 5D and *QPms.inra-6A2* located on chromosome 6A (Muranty et al. 2008). Two major quantitative trait loci (QTL), *QPm.inra-5D* and *QPm.inra-6A2*, control RE714 powdery mildew resistance at the adult stage and were consistently detected in the studies of Chantret et al. (2001), Mingeot et al. (2002) and Muranty et al. (2009), involving crosses of RE714 with the susceptible lines 'Hardi' or 'Festin'. A third region involved in powdery mildew resistance at the adult stage was detected on chromosome 7A by Chantret et al. (2001) and by Muranty et al. (2009), who described two closely located QTL expressed in different years. This region had a rather minor effect (Table 1). The genetic factors involved in resistance at the seedling stage and at the adult stage on chromosome 5D and 6A are located in the same regions. This co-localisation probably explains what was originally called the residual effect of *MIRE* (Chantret et al. 1999): *MIRE* was then a resistance gene of RE714 defined by resistance scoring at the seedling stage with specific isolates ('93-27' and '95-9') in RE714 x Hardi segregating populations, but this resistance was later broken up into the two QTL, *QPms.inra-5D* and *QPms.inra-6A2*, and *MIRE* was re-defined as the resistance factor located on chromosome 6A (Chantret et al. 2000). The localization and effects of these QTL are detailed in Table 1.

To assess the origin of the main genetic factors involved in RE714 powdery mildew resistance, the linked microsatellite markers obtained by Muranty et al. (2008; 2009) were used to compare RE714 alleles to the alleles of its ancestral lines.

## Materials and methods

### Plant material

The pedigree of RE714 includes two interspecific crosses and several crosses with known wheat cultivars, except for the unknown genotype that pollinated the single male-sterile plant obtained from the cross *Ae. tauschii* no. 33 × *T. dicoccum* no. 119 had a progeny by open

pollination (Fig. 1). The wheat lines VPM, 'Marne', 'Moisson' and 'Beauchamp', and the lines *Ae. tauschii* n°33, *Ae. ventricosa* n°10 and *T. dicoccum* n°119 were used to infer the origins of the main genetic factors involved in RE714 powdery mildew resistance. Unfortunately, the *T. persicum* accession that was used to develop VPM has been lost.

Only *T. dicoccum* n°119 was tested in the seedling stage resistance tests described by Muranty et al. (2008) because it was used as a differential check. *T. dicoccum* n°119 is resistant to specific isolates of powdery mildew at the seedling stage (isolates 'K', '93-27', '95-9', '95-31' and '97-23' in Muranty et al. 2008, 15 isolates in Robe and Doussinault 1995) and RE714 is also resistant to these isolates (Muranty et al. 2008, Robe and Doussinault 1995). *T. dicoccum* n°119 was described as the donor of *MIRE* by Robe and Doussinault (1995). 'Beauchamp' and 'Moisson' were reported to be susceptible to 12 differential powdery mildew isolates at the seedling stage by Zeller et al. (1993) and to carry no known resistance gene. 'Marne' was susceptible to a local powdery mildew isolate in Australia and is not known to carry *Pm4b* (Bariana and McIntosh 1994). VPM is resistant to specific isolates of the powdery mildew pathogen at the seedling stage and carries *Pm4b* (Bariana and McIntosh 1993, Zeller et al. 1993). *Ae. tauschii* n°33 was resistant to all 24 isolates tested by Robe and Doussinault (1995) at the seedling stage. VPM, 'Beauchamp', *Ae. tauschii* n°33 and *T. dicoccum* n°119 were tested in the adult plant resistance tests described by Muranty et al (2009). Area under disease progress curve (AUDPC) were calculated using several disease assessments as  $AUDPC = \sum_i [x_i + x_{i+1}] \frac{t_{i+1} - t_i}{2}$  where  $x_i$  is the score at date  $t_i$ . The mean AUDPC of 'Hardi', RE714 and its tested parental lines in each test year are reported in Table 2. A multiple mean comparison test was performed with the TukeyHSD function under R (R-Development-Core-Team 2006) to identify significant differences between tested lines.

## Microsatellite marker analysis

Genomic DNA was extracted from fresh leaves following a modified CTAB method (Doyle and Doyle 1990). The DNA concentrations were adjusted to 10 ng/μl. PCR reactions for microsatellite markers were performed as described by Tixier et al. (1997). PCR products were separated on 6% polyacrylamide denaturing gels and visualized by silver nitrate staining or using a Li-Cor DNA analyzer (Li-Cor Inc., Lincoln, NE). For the latter system, each right primer was 5'-tailed with the M13 forward consensus sequence. The M13-tailed right primers were then used in combination with a standard M13 primer dye-labelled at the 5'-end (Boutin-Ganache et al. 2001). A total of 36 wheat microsatellite primer pairs were used to genotype

the lines. These markers were chosen from the map of the RE714 x Hardi RIL population described in Muranty et al. (2009) to have a good representation of the support intervals (-1.5 LOD) of the main powdery mildew resistance QTL detected in this population. The data were recorded as "a band of the same size as the band of RE714" (RE), "a band of the same size as the band of 'Hardi'" (Ha), "a band with a size different from the sizes of the bands of RE714 and 'Hardi'" (D) or "no band" (NO) (Table 3).

## Results

### Adult plant reactions of some of the RE714 ancestral lines

Adult plant actions of VPM, 'Beauchamp', *Ae. tauschii* n°33 and *T. dicoccum* n°119 were evaluated together with the RE714 x Hardi RIL population (Muranty et al. 2009). Mean AUDPC for powdery mildew were generally higher in 2001 and 2002 when the tests were performed in tunnel houses which provided an environment more favourable to powdery mildew development because they were warmer and more humid (the plants were regularly watered) than the field. Three and four groups of lines similar in AUDPC means were identified with the multiple mean comparison test in 2000 and 2001, respectively (Table 2). *Ae. tauschii* no. 33 was highly resistant at the adult plant stage in the 3 years, and its mean AUDPC was lower than or very close to the RE714 mean AUDPC: the differences between *Ae. tauschii* no. 33 and RE714 were not significant ( $P = 0.05$ ). *T. dicoccum* n°119 had a mean AUDPC very close to RE714 mean AUDPC in 2000, but was significantly more susceptible than RE714 in 2001. In 2002, it had also an intermediate response level. 'Beauchamp' had an intermediate response level in all 3 years, which resulted in a mean AUDPC below or near the mean of the recombinant inbred lines from the cross RE714 x 'Hardi'. However, 'Beauchamp' was not significantly different from RE714 in 2000 whereas it was significantly more susceptible than RE714 in 2001. 'Beauchamp' was significantly more resistant than 'Hardi' in 2000 and 2001. These results confirmed the weak adult plant resistance of 'Beauchamp' that was recorded when it was registered in (Anonymous 1979). VPM was susceptible in all 3 years and was not significantly different from 'Hardi' in 2000 and 2001. It was also not significantly different from 'Beauchamp' in 2000.

The origin of the main QTL was investigated by comparing the microsatellite marker alleles of RE714 to those of its parental lines. These markers had been mapped in a RE714 x Hardi RIL population (Muranty et al. 2009) (see Fig. 2). For each genetic factor, we focused our attention to the parental lines that could be donors of the QTL, and excluded the lines that did not possess the resistance factor carrying genome. Indeed, *Ae. tauschii* n°33 (DD) and *Ae*

*ventricosa* n°10 (DDN<sup>V</sup>N<sup>V</sup>) as well as *T. dicoccum* n°119 (AABB) cannot be the donors of genetic factors located in the A and D genomes, respectively.

## Origin of *Pm4b*, a gene mapped on chromosome 2A

Because 'Beauchamp', 'Marne' and 'Moisson' were recorded as susceptible to all tested powdery mildew isolates at the seedling stage, including isolates avirulent on *Pm4b* reference lines, they cannot be the donor of *Pm4b*. *T. dicoccum* n°119 is susceptible to some isolates that are avirulent on reference lines carrying *Pm4b* (Robe and Doussinault, 1995; Muranty et al 2008) so it can not be the donor of *Pm4b*. This leaves only VPM and the unknown line that pollinated the single male-sterile plant obtained in the initial cross between *Ae. tauschii* no. 33 and *T. dicoccum* no. 119 as potential donor of *Pm4b* to RE714. The microsatellite loci located within the support interval of *QPms.inra-2A3*, corresponding to the *Pm4b* gene (Muranty et al 2008), are *Xgwm311*, *Xgpw4474* and *Xgpw4456* (Fig. 2). In the vicinity of the *Pm4b* gene on linkage group 2A3, RE714 had the same alleles as VPM at microsatellite loci *Xcfd0251*, *Xgwm526* and *Xgwm311*, but not at the *Xgwm382* and *Xgpw4474* loci (Table 3). This confirmed that VPM is the most likely donor of *Pm4b* to RE714. Among the parental lines of VPM, the lines that contributed to the A genome are 'Marne' and the *T. persicum* line, which is the most likely donor of *Pm4b* because 'Marne' is susceptible.

## Origin of *QPm.inra-5D*

The microsatellite loci located within the support interval of *QPm.inra-5D* are *Xcfd0008*, *Xgwm494-5D*, *Xgwm639-5D*, *Xgpw5207-5D*, *Xcfd0057* and *Xcfd0026* (Fig. 2). The putative wild species donors of this resistance factor are *Ae. tauschii* n°33 (DD) and *Ae ventricosa* n°10 (DDN<sup>V</sup>N<sup>V</sup>). RE714 had the same allele as *Ae. tauschii* n°33, which is completely resistant to powdery mildew, at all six microsatellite loci located within the support interval of *QPm.inra-5D*, but not at other nearby loci located outside the support interval, namely *Xgpw4020*, *Xgwm159*, *Xcfd0007*, *Xcfd0003* and *Xcfd0012*. On the contrary, *Ae ventricosa* n°10 had an allele different from the RE714 allele at four of the six microsatellite loci located within the support interval of *Qpm.inra-5D*. Consequently, the most probable origin of *QPm.inra-5D* in RE714 was *Ae. tauschii* n°33. The QTL *QPm.inra-5D* explained up to 66.7% of the phenotypic variance at the seedling stage and up to 56.3% at the adult plant stage (Table 1), but its additive effects (-22.1 for AUDPC in 2000, -68.6 in 2001 and -46.0 in 2002, see Table 3 in Muranty et al (2009)) are not high enough to explain the difference between *Ae. tauschii* n°33 and the mean of the RE714 x Hardi RIL population (Table 2). Besides, the

results indicate an interstitial position of the fragment of *Ae. tauschii* n°33 introgressed in the 5D chromosome of RE714. The length of this introgressed fragment was approximately 40 cM on the map obtained in the RE714 x Hardi RIL population (Muranty et al 2009).

## Origin of *QPm.inra-6A2*

The microsatellite loci located within the support interval of *QPm.inra-6A2* and *QPms.inra-6A2a* are *Xgpw7388* and *XDUPw167*, whereas only one microsatellite locus, namely *Xgwm427*, was located within the support interval of *QPms.inra-6A2b*. The putative wild species donors of resistance factors in this region are *T. dicoccum* n°119 (AABB) and the lost *T. persicum* accession that was used to develop VPM. *T. dicoccum* n°119 was previously suggested as the donor of the so-called *MIRE* resistance (Robe and Doussinault 1995). RE714 had the same allele as *T. dicoccum* n°119 at microsatellite loci *Xgwm617*, *Xcfa2114*, *Xgwm427*, *Xbarc104*, *Xgpw7388* and *XDUPw167*, but not at *Xgwm169*, whereas VPM had the same allele as RE714 at only one of these loci, namely *Xbarc104*. We could thus confirm that the most probable origin of *QPm.inra-6A2* in RE714 was *T. dicoccum* n°119, as already suggested by Robe and Doussinault (1995) and Chantret et al. (2000), and show that the fragment of *T. dicoccum* n°119 introgressed on the 6A chromosome of RE714 has a distal position. The length of this introgressed fragment was approximately 76 cM on the map obtained in the RE714 x Hardi RIL population (Muranty et al 2009).

## Origin of *QPm.inra-7A3a* and *QPm.inra-7A3b*

The microsatellite loci located within the support interval of *QPm.inra-7A3a* are *Xgpw2308*, *Xgpw2338*, *Xgpw2299*, *Xcfa2040a* and *Xgwm146*. This QTL could correspond to the QTL detected by Chantret et al (2001) near *Xgwm344* in an F<sub>2:3</sub> population derived from the cross RE714 x Hardi. RE714 had the same allele as 'Beauchamp' and 'Moisson' at the microsatellite loci *Xgpw2338* and *Xgpw2299*, and also the same as 'Moisson' at *Xgwm146*, but not at the other loci. 'Marne' had the same allele as RE714 at the microsatellite loci *Xgpw2338* and *Xgwm146* and not at the other loci but VPM had the same allele as RE714 only at the microsatellite locus *Xgpw2338*. The accession *T. dicoccum* n°119 also had the same allele as RE714 at the microsatellite locus *Xgpw2299* and *Xgwm344*. Consequently, it was difficult to identify the donor of *QPm.inra-7A3a* among the parental line of RE714 that were tested. The unknown line that pollinated the single male-sterile plant obtained in the initial cross between *Ae. tauschii* no. 33 and *T. dicoccum* no. 119 could be the donor.



The microsatellite loci located within the support interval of *QPm.inra-7A3b* are *Xgpw2252* and *Xcfa2019*. 'Beauchamp' was the only tested parental line to have the same allele as RE714 at these loci, whereas 'Moisson' had the same allele as RE714 only at *Xcfa2019*. 'Beauchamp' is susceptible to all powdery mildew isolates at the seedling stage, but has an intermediate response level at the adult stage (Table 2, and see above). As the QTL *QPm.inra-7A3b* is expressed only at the adult stage and its effect is quite weak (Table 1), 'Beauchamp' could be the origin of the favourable allele in RE714.

## Discussion

Several powdery mildew resistance genes have been introduced from bread wheat related species with an AABB genome: *Pm3h* and *Mld* from *T. durum*, *Pm4a* and *Pm5a* from *T. dicoccum*, *Pm4b*, *Pm33* and *PmPs5A* from *T. carthlicum*, and *Pm16*, *Pm26*, *Pm30*, *Pm31*, *MITd1055* and *Mlze1* from *T. dicoccoides* (McIntosh et al. 2008). In RE714, *Pm4b* probably came from *T. persicum* = *T. carthlicum* through VPM, so was introduced into bread wheat differently from the *Pm4b* gene in 'Armada', which has been the reference line for *Pm4b* in several studies (Heun and Fischbeck 1987a,b, Singrun et al. 2004). RE714 QTL on chromosome 6A is another interesting gene probably introduced from one of these species, namely *T. dicoccum*. This illustrates the value of these species as genetic resources for the improvement of resistance to biotic stresses in bread wheat.

*Ae. tauschii* was apparently not a common source of powdery mildew resistance genes until recently. Among the named powdery mildew resistance genes, *Pm2*, *Pm19*, *Pm34* and *Pm35* were introduced from this species into wheat. However, *Pm2* is widespread in cultivars grown in Europe and China (Hsam and Zeller 2002) and its origin is often more probably in old wheat varieties or landraces than in an introgression from *Ae. tauschii*. Within the CIMMYT wheat breeding programs, hundreds of 'synthetic' wheat lines, i.e. hybrids between species with an AABB genome and *Ae. tauschii*, have been created and their derivatives now represent a significant proportion of the parental lines used in crosses in these breeding programmes each year (Crouch et al 2009). These lines may contain additional powdery mildew resistance genes. Our results concerning common alleles in RE714 and *Ae. tauschii* n°33 strongly suggest that the RE714 QTL on chromosome 5D originated in this accession. However, the resistance to powdery mildew conferred by this QTL is not as strong as the resistance found in *Ae. tauschii* n°33, either because *Ae. tauschii* n°33 carries other resistance genes, or because the gene does not have the same level of expression in the hexaploid background as in the diploid. *Ae. tauschii* n°33 was one of the parents of synthetics studied by

Lutz et al. (1994) who showed that synthetic amphiploids generally express lower levels of resistance to powdery mildew than the respective diploid parental lines.

Reduced recombination is perceived as a general drawback of introgressions from related species. For example, in the vicinity of *Pchl1*, a gene for eyespot resistance introduced from *Ae. ventricosa* into wheat through VPM, 6 markers showed complete linkage in a population of 254 RIL segregating for *Pchl1*, whereas three of these markers spanned a 3.8 cM segment in the ITMI population that does not segregate for *Pchl1* (Leonard et al. 2008). On the contrary, fragments originating from *Ae. tauschii* n°33 and *T. dicoccum* n°119 around *QPm.inra-5D* and *QPm.inra-6A2* in RE714 did not seem to have reduced recombination. Consequently, developing advanced intercross lines with RE714 could help delineate its powdery mildew resistance factors to smaller intervals and perhaps result in genotypes with reduced linkage drag.

More generally, wheat breeders should enlarge their use of related species from the primary gene pool, for example by creating and exploiting 'synthetic wheat', because genetic diversity within these species does not seem to be fully exploited and introgression of interesting genetic factors from these species does not require difficult chromosome engineering strategies. Often, the value of such genotypes for an agronomic trait will not be predictive of the value of the genotypes obtained from them in synthetic wheats or direct hybrid derivatives, as seen in the present example of powdery mildew resistance in RE714 and *Ae. tauschii* n°33. A strategy to introduce genetic diversity from related species of the primary gene pool into wheat would be to perform widescale advanced backcrosses, by creating intercross populations involving several synthetic wheat genotypes and modern breeding lines and then apply recurrent selection for adaptation to modern cultivation techniques and desirable agronomic traits, with mild selection pressure in order to avoid a too rapid shrinkage of genetic diversity.

## Acknowledgements

Molecular mapping results were in great part obtained through the OUEST-genopole® genotyping technical platform, successively supervised by S. Prioul, A. Lostanlen and S. Coedel. We thank P. Sourdille for sharing information about microsatellite primer sequences ahead of publication. We thank J. Jahier and M. Trottet for their valuable comments on the first draft of this article. We thank the reviewers for their careful reading which helped us improve this paper.

## References

- Anonymous, 1979: Bulletin des variétés. Céréales & fiches descriptives des variétés inscrites en octobre 1978. GEVES, La Minière, France.
- Bariana, H.S., and R.A. McIntosh, 1993: Cytogenetic studies in wheat. XV. Location of rust resistance genes in VPM1 and their genetic linkage with other disease resistance genes in chromosome 2A. *Genome* **36**, 476-482.
- Bariana, H.S., and R.A. McIntosh, 1994: Characterisation and origin of rust and powdery mildew resistance genes in VPM1 wheat. *Euphytica* **76**, 53-61.
- Boutin-Ganache, I., M. Raposo, M. Raymond, and C.F. Deschepper, 2001: M13-tailed primers improve the readability and usability of microsatellite analyses performed with two different allele-sizing methods. *BioTechniques* **31**, 24-27.
- Chantret, N., M.T. Pavoine, and G. Doussinault, 1999: The race-specific resistance gene to powdery mildew, *MIRE*, has a residual effect on adult plant resistance of winter wheat line RE714. *Phytopathology* **89**, 533-539.
- Chantret, N., P. Sourdille, M. Roder, M. Tavaud, M. Bernard, and G. Doussinault, 2000: Location and mapping of the powdery mildew resistance gene *MIRE* and detection of a resistance QTL by bulked segregant analysis (BSA) with microsatellites in wheat. *Theor. Appl. Genet.* **100**, 1217-1224.
- Chantret, N., D. Mingeot, P. Sourdille, M. Bernard, J.M. Jacquemin, and G. Doussinault, 2001: A major QTL for powdery mildew resistance is stable over time and at two development stages in winter wheat. *Theor. Appl. Genet.* **103**, 962-971.
- Crouch, J.H., T.S. Payne, S. Dreisigacker, H. Wu, and H.-J. Braun, 2009: Improved Discovery and Utilization of New Traits for Breeding Wheat Facts and Futures 2009, 42-51. CIMMYT, Mexico, D.F.
- Doyle, J.J., and J.L. Doyle, 1990: Isolation of DNA from fresh tissue. *Focus* **12**, 13-15.
- Feuillet, C., P. Langridge, and R. Waugh, 2008: Cereal breeding takes a walk on the wild side. *Trends Genet.* **24**, 24-32.
- Heun, M., and G. Fischbeck, 1987a: Genes for powdery mildew resistance in cultivars of spring wheat. *Plant Breeding* **99**, 282-288.
- Heun, M., and G. Fischbeck, 1987b: Identification of wheat powdery mildew resistance genes by analysing host-pathogen interactions. *Plant Breeding* **98**, 124-129.
- Hospital, F., 2005: Selection in backcross programmes. *Philos. Trans. R. Soc. Lond., Ser. B: Biol. Sci.* **360**, 1503-1511.

# Postprint

Version définitive du manuscrit publié dans / Final version of the manuscript published in :  
Plant breeding, 2010, 129 (5), 465-471 <http://dx.doi.org/10.1111/j.1439-0523.2009.01740.x>

Hsam, S.L.K., and F.J. Zeller, 2002: Breeding for powdery mildew resistance in common wheat (*Triticum aestivum* L.), In: R. R. Bélanger, W. R. Bushnell, A. J. Dik and T. L. W. Carver, (eds.) Powdery Mildews: a Comprehensive Treatise, 219-238. Amer Phytopathological Soc, St Paul.

Jiang, J.M., B. Friebe, and B.S. Gill, 1994: Recent advances in alien gene-transfer in wheat. *Euphytica* **73**, 199-212.

Leonard, J., C. Watson, A. Carter, J. Hansen, R. Zemetra, D. Santra, K. Campbell, and O. Riera-Lizarazu, 2008: Identification of a candidate gene for the wheat endopeptidase *Ep-D1* locus and two other STS markers linked to the eyespot resistance gene *Pch1*. *Theor. Appl. Genet.* **116**, 261-270.

Lutz, J., S.L.K. Hsam, E. Limpert, and F.J. Zeller, 1994: Powdery mildew resistance in *Aegilops tauschii* Coss. and synthetic hexaploid wheats. *Genet. Resour. Crop Evol.* **41**, 151-158.

McIntosh, R.A., Y. Yamazaki, J. Dubcovsky, J. Rogers, C. Morris, D.J. Somers, R. Appels, and K.M. Devos, 2008: Catalogue of Gene Symbols for Wheat.

Mingeot, D., N. Chantret, P.V. Baret, A. Dekeyser, N. Boukhatem, P. Sourdille, G. Doussinault, and J.M. Jacquemin, 2002: Mapping QTL involved in adult plant resistance to powdery mildew in the winter wheat line RE714 in two susceptible genetic backgrounds. *Plant Breeding* **121**, 133-140.

Muranty, H., M.T. Pavoine, B. Jaudeau, W. Radek, G. Doussinault, and D. Barloy, 2008: A quantitative approach detects three QTLs involved in powdery mildew resistance at the seedling stage in the winter wheat line RE714. *Aust. J. Agric. Res.* **59**, 714-722.

Muranty, H., M.T. Pavoine, B. Jaudeau, W. Radek, G. Doussinault, and D. Barloy, 2009: Two stable QTL involved in adult plant resistance to powdery mildew in the winter wheat line RE714 are expressed at different times along the growing season. *Mol. Breed.* **23**, 445-461.

Qi, L.L., B. Friebe, P. Zhang, and B.S. Gill, 2007: Homoeologous recombination, chromosome engineering and crop improvement. *Chromosome Res.* **15**, 3-19.

R-Development-Core-Team, 2006: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Robe, P., and G. Doussinault, 1995: Genetic analysis of powdery-mildew resistance of a winter-wheat line, RE714, and identification of a new specific-resistance gene. *Plant Breeding* **114**, 387-391.

Comment citer ce document :

Muranty, H., Pavoine, M.-T., Doussinault, G., Barloy, D. (2010). Origin of powdery mildew resistance factors in RE714, a wheat breeding line obtained from two interspecific crosses. *Plant Breeding*, 129 (5), 465-471. DOI : 10.1111/j.1439-0523.2009.01740.x

# Postprint

Version définitive du manuscrit publié dans / Final version of the manuscript published in :  
Plant breeding, 2010, 129 (5), 465-471 <http://dx.doi.org/10.1111/j.1439-0523.2009.01740.x>

Roussel, V., J. Koenig, M. Beckert, and F. Balfourier, 2004: Molecular diversity in French bread wheat accessions related to temporal trends and breeding programmes. *Theor. Appl. Genet.* **108**, 920-930.

Singrun, C., S.L.K. Hsam, F.J. Zeller, G. Wenzel, and V. Mohler, 2004: Localization of a novel recessive powdery mildew resistance gene from common wheat line RD30 in the terminal region of chromosome 7AL. *Theor. Appl. Genet.* **109**, 210-214.

Tixier, M.H., P. Sourdille, M. Röder, P. Leroy, and M. Bernard, 1997: Detection of wheat microsatellites using a non radioactive silver-nitrate staining method. *J. Genet. Breed.* **51**, 175-177.

Trottet, M., and F. Dosba, 1983: Cytogenetical analysis and behavior towards *Septoria nodorum* of hybrids between *Triticum* sp and *Aegilops squarrosa* and of their progenies. *Agronomie* **3**, 659-664.

Xie, W.L., and E. Nevo, 2008: Wild emmer: genetic resources, gene mapping and potential for wheat improvement. *Euphytica* **164**, 603-614.

Zeller, F.J., J. Lutz, E.I. Reimlein, E. Limpert, and J. Koenig, 1993: Identification of powdery mildew resistance genes in common wheat (*Triticum aestivum* L). II. French cultivars. *Agronomie* **13**, 201-207.

**Table 1** Main genetic factors controlling RE714 powdery mildew resistance combining data at the seedling stage (15 isolates in a RIL population, three isolates in an F<sub>2:3</sub> population) and at the adult plant stage (three years in a RIL and a DH population, 2 years in an F<sub>2:3</sub> population)

Chr. region	QTL	Resistance factor		QTL characteristics	Upstream flanking marker	Peak position		Ref.
		Growth stage	Population			(cM) <sup>1</sup>	R <sup>2</sup> % <sup>2</sup>	
2A3	<i>QPms.inra-2A3</i>	seedling	RIL	6 isolates	gpw4456	0.0 – 2.0	76.0 – 92.9	Muranty et al. (2008)
5D	<i>QPms.inra-5D</i>	seedling	RIL	12 isolates	cf0026 (9 isolates) or gpw5207-5D (2 isolates) or gwm494-5D (1 isolate)	0.0 – 8.0	10.6 – 66.7	Muranty et al. (2008)
			F <sub>2:3</sub>	3 isolates	gwm174	-	16.8 – 25.3	Chantret et al. (2000)
	<i>QPm.inra-5D</i>	adult	RIL	AUDPC (3 years) and individual scores (4 or 5 each year)	cf0026 (15 traits) or gpw5207-5D (1 trait)	0.0 – 2.0	8.5 – 56.3	Muranty et al. (2009)
			DH	AUDPC (3 years)	cf0026 (1 year) or gbxG083c (2 years)	0.0 – 4.0	33.5 – 37.9	Chantret et al. (2001)
			F <sub>2:3</sub>	AUDPC (2 years)	cf0026	2.0	28.1 – 37.4	Chantret et al. (2001)
6A2	<i>QPms.inra-6A2a</i>	seedling	RIL	2 isolates	gpw7388	4.0	19.8 – 30.7	Muranty et al. (2008)
			F <sub>2:3</sub>	3 isolates	ksuD27	-	24.1 – 37.0	Chantret et al. (2000)
	<i>QPms.inra-6A2b</i>	seedling	RIL	1 isolate	gwm427	0.0	14.1	Muranty et al. (2008)
	<i>QPm.inra-6A2</i>	adult	RIL	AUDPC (1 year) and individual scores (1 to 4 each year)	gpw7388	2.0 – 4.0	12.2	Muranty et al. (2009)
			DH	AUDPC (1 year)	MIRE	-	12.2	Chantret et al. (2001)
			F <sub>2:3</sub>	AUDPC (2 years)	gwm427	6.0 – 8.0	8.8 – 13.4	Chantret et al. (2001)
7A3	<i>QPm.inra-7A3-a</i>	adult	RIL	2 individual scores in 2000	P32M51n	6.0	10.8 – 18.7	Muranty et al. (2009)
			F <sub>2:3</sub>	AUDPC (1 year)	gwm344	4.0	6.4	Chantret et al. (2001)
	<i>QPm.inra-7A3-b</i>	adult	RIL	AUDPC in 2001	gpw2252	8.0	5.9	Muranty et al. (2009)

<sup>1</sup> Distance to the upstream flanking marker in the map obtained in the RE714 x 'Hardi' tested populations: RIL (Muranty et al. 2008), HD or F<sub>2:3</sub> (Chantret et al. 2001).

<sup>2</sup> Percentage of phenotypic variance explained.

# Postprint

Version définitive du manuscrit publié dans / Final version of the manuscript published in :  
Plant breeding, 2010, 129 (5), 465-471 <http://dx.doi.org/10.1111/j.1439-0523.2009.01740.x>

Table 2: Mean AUDPC for powdery mildew responses of RE714, 'Hardi', some of the parental lines of RE714 and the RIL derived from the cross RE714 x 'Hardi', in adult stage reaction tests described by Muranty et al. (2009). These tests were performed under naturally infected field conditions in Le Rheu (France), in the field in 2000 and in tunnel houses in 2001 and 2002. The 2000 and 2001 tests were sown in October in 1999 and 2000, respectively, whereas the 2002 test was sown in February 2002.<sup>1</sup>

Year	RE714	Hardi	<i>Ae. tau</i> n° 33	<i>T. dic</i> n° 119	Beauchamp	VPM	RIL
2000	84.5a	238.5c	87.5a	87.5a	116.7ab	184.5bc	149
2001	325ab	668d	280a	481c	438bc	601d	463
2002 <sup>2</sup>	170	543	58	254	320	400	312

<sup>1</sup> Mean AUDPC followed by different letters are significantly different.

<sup>2</sup> Statistical assessment of mean differences could not be performed with the 2002 data.

**Table 3** Genotypes of RE714 and its parental lines and of 'Hardi' at microsatellite loci found in the vicinity of RE714 QTL for powdery mildew resistance.

Chr.	region	marker	RE714	Hardi	<i>Aegilops ventricosa</i>		<i>Aegilops tauschii</i>		<i>Triticum dicoccum</i>		
					VPM	no. 10	no. 33	n°119	Beauchamp	Marne	Moisson
2A3		cfD0251	RE	Ha	RE	Ha	D	D	Ha	RE?	Ha
		gwm526	NO	Ha	NO	D	Ha	NO	NO	NO	Ha
		gwm382	RE	NO	D	D	D	D	RE	D	RE
		<b>gwm311</b>	<b>RE</b>	<b>NO</b>	<b>RE</b>	<b>D</b>	<b>D</b>	<b>D</b>	<b>RE</b>	<b>NO</b>	<b>RE</b>
		<b>gpw4474</b>	<b>NO</b>	<b>-</b>	<b>D</b>	<b>Ha</b>	<b>NO</b>	<b>Ha?</b>	<b>NO</b>	<b>-</b>	<b>-</b>
5D		gpw4020	RE	Ha	Ha?	D	D	D	RE	Ha	RE
		gwm159	RE	Ha	Ha	D	D	-	Ha	Ha	Ha
		<b>cfD0008</b>	<b>RE</b>	<b>Ha</b>	<b>Ha?</b>	<b>D</b>	<b>RE</b>	<b>D</b>	<b>D</b>	<b>-</b>	<b>D</b>
		<b>gwm494-5D</b>	<b>NO</b>	<b>Ha</b>	<b>Ha</b>	<b>D</b>	<b>NO</b>	<b>NO</b>	<b>Ha</b>	<b>Ha</b>	<b>Ha</b>
		<b>gwm639-5D</b>	<b>RE</b>	<b>Ha</b>	<b>Ha</b>	<b>RE?</b>	<b>RE</b>	<b>-</b>	<b>Ha</b>	<b>Ha</b>	<b>Ha</b>
		<b>gpw5207-5D</b>	<b>RE</b>	<b>Ha</b>	<b>-</b>	<b>-</b>	<b>RE</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>-</b>
		<b>cfD0057</b>	<b>RE</b>	<b>Ha</b>	<b>D</b>	<b>D</b>	<b>RE</b>	<b>-</b>	<b>Ha</b>	<b>D</b>	<b>D</b>
		<b>cfD0026</b>	<b>RE</b>	<b>Ha</b>	<b>D</b>	<b>D</b>	<b>RE</b>	<b>D</b>	<b>Ha</b>	<b>D</b>	<b>D</b>
		cfD0007	RE	Ha	RE	D	D	-	RE	-	RE
		cfD0003	RE	Ha	D	D	D	-	RE	RE	RE
6A2		cfD0012	RE	Ha	RE?	-	D	-	RE	RE	RE
		gwm169	RE	Ha	-	-	-	D	D	RE	RE
		gwm617	RE	Ha	D	-	-	RE	-	-	-
		cfa2114	RE	Ha	D	-	-	RE	D	Ha	Ha
		<b>gwm427</b>	<b>RE</b>	<b>Ha</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>RE</b>	<b>D</b>	<b>-</b>	<b>-</b>
		barc104	NO	Ha?	NO	NO	Ha	NO	D	D	D
		<b>gpw7388</b>	<b>NO</b>	<b>Ha?</b>	<b>Ha</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>Ha</b>	<b>Ha</b>	<b>Ha</b>
7A3		<b>DuPw167</b>	<b>RE</b>	<b>Ha</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>RE</b>	<b>-</b>	<b>Ha</b>	<b>Ha</b>
		gpw2119	RE	Ha	Ha	-	-	D	RE	Ha	Ha
		gpw2264	RE	Ha	Ha	-	-	Ha	RE	Ha	Ha
		gpw2103	RE	Ha	-	-	-	D	RE	-	Ha
		barc121	RE	Ha	-	-	D	RE	D	-	D
		gwm282	RE	Ha	D	-	-	D	-	D	D
		<b>gpw2252</b>	<b>RE</b>	<b>Ha</b>	<b>Ha</b>	<b>-</b>	<b>-</b>	<b>Ha</b>	<b>RE</b>	<b>Ha</b>	<b>Ha</b>
		<b>cfa2019</b>	<b>RE</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>D</b>	<b>RE</b>	<b>NO</b>	<b>RE</b>
		<b>gpw2308</b>	<b>RE</b>	<b>Ha</b>	<b>D</b>	<b>-</b>	<b>-</b>	<b>D (Ha)</b>	<b>D</b>	<b>D</b>	<b>D</b>
		<b>gpw2338</b>	<b>RE</b>	<b>Ha</b>	<b>RE</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>RE</b>	<b>RE</b>	<b>RE</b>
		<b>gpw2299</b>	<b>RE</b>	<b>Ha</b>	<b>Ha?</b>	<b>-</b>	<b>-</b>	<b>RE</b>	<b>RE</b>	<b>-</b>	<b>RE</b>

Commenter ce document :



<b>cfa2040a</b>	<b>RE</b>	<b>Ha</b>	-	-	-	<b>Ha</b>	-	-	-
<b>gwm146</b>	<b>RE</b>	<b>Ha</b>	<b>D</b>	-	-	<b>D</b>	<b>D</b>	<b>RE</b>	<b>RE</b>
gwm344	RE	Ha	D	-	-	RE	Ha	D	Ha

Loci located within support intervals of QTL are in bold.

<sup>1</sup> Genotypes were scored with reference to the genotypes of RE714 and 'Hardi'

RE, a band of the same size as the band of RE714; Ha, a band of the same size as the band of 'Hardi'; RE?, Ha?, a band of the same size as the band of RE714, or 'Hardi', but much fainter or much stronger; D, a band with a molecular weight different from those of RE714 and 'Hardi'; NO, no band; -, no amplification when the marker is codominant in RE714 x 'Hardi'.

**Supplementary table 1** Band sizes of RE714 and its parental lines and of 'Hardi' at microsatellite loci found in the vicinity of RE714 QTL for powdery mildew resistance. Loci located within support intervals of QTL are in bold.

Chr. region	Marker	RE714	Hardi	VPM	<i>Ae. vent.</i> n°10	<i>Ae. tau.</i> n°33	<i>T. dic.</i> n°119	Beauchamp	Marne	Moisson
2A3	cfD0251									
	gwm526	NO		NO			NO	NO	NO	
	gwm382	133	NO	152	139/165	139	197	133	152	133
	<b>gwm311</b>		<b>NO</b>						<b>NO</b>	
5D	<b>gpw4474</b>	<b>NO</b>	-		<b>157</b>	<b>NO</b>	<b>157?</b>	<b>NO</b>	-	-
	gpw4020	184	186	186	190	178	190	184	186	184
	gwm159						-			
	<b>cfD0008</b>	<b>171</b>	<b>175</b>	<b>175?</b>	<b>161</b>	<b>171</b>	<b>161</b>	<b>173</b>	-	<b>173</b>
	<b>gwm494-5D</b>	<b>NO</b>	<b>197</b>	<b>197</b>	<b>191</b>	<b>NO</b>	<b>NO</b>	<b>197</b>	<b>197</b>	<b>197</b>
	<b>gwm639-5D</b>	<b>138</b>	<b>154</b>	<b>154</b>	<b>138?</b>	<b>138</b>	-	<b>154</b>	<b>154</b>	<b>154</b>
	<b>gpw5207-5D</b>	<b>207</b>	<b>193</b>	-	-	<b>207</b>	-	-	-	-
	<b>cfD0057</b>	<b>332</b>	<b>294</b>	<b>300</b>	<b>275</b>	<b>332</b>	-	<b>294</b>	<b>300</b>	<b>300</b>
	<b>cfD0026</b>									
	cfD0007	259	255	259		274	-	259	-	259
6A2	cfD0003	194	190	192	192	184	-	194	194	194
	cfD0012	210	208	210	-	214	-	210	210	210
	gwm169	216	240	-	-	-	214	214	216	216
	gwm617							-	-	-
	cfa2114									
	<b>gwm427</b>	<b>216</b>	<b>228</b>	-	-	-	<b>216</b>	<b>240</b>	-	-
	barc104	NO	228	NO	NO	228	NO			
	<b>gpw7388</b>	<b>NO</b>	<b>123</b>	<b>123</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>123</b>	<b>123</b>	<b>123</b>
	<b>DuPw167</b>	<b>244</b>	<b>256</b>	-	-	-	<b>244</b>	-	<b>256</b>	<b>256</b>
	gpw2119	220	218	218	-	-	202	220	218	218
7A3	gpw2264									
	gpw2103	255	263	-	-	-	245	255	-	263
	barc121	246	235	-	-	-	250	245	-	235
	gwm282									
	<b>gpw2252</b>	<b>259</b>	<b>253</b>	<b>253</b>	-	-	<b>253</b>	<b>259</b>	<b>253</b>	<b>253</b>
	<b>cfa2019</b>	<b>240</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>	<b>246</b>	<b>240</b>	<b>NO</b>	<b>240</b>
	<b>gpw2308</b>	<b>260</b>	<b>200</b>	<b>256</b>	-	-	<b>200/186</b>	<b>258</b>	<b>254</b>	<b>258</b>
	<b>gpw2338</b>	<b>180</b>	<b>190</b>	<b>180</b>	-	-	-	<b>180</b>	<b>180</b>	<b>180</b>
	<b>gpw2299</b>	<b>272</b>	<b>266</b>	<b>266</b>	-	-	<b>272</b>	<b>272</b>	-	<b>272</b>
	<b>cfa2040a</b>	<b>333</b>	<b>313</b>	-	-	-	<b>313</b>	-	-	-

Commenter ce document :

**gwm146**  
**gwm344**

# Postprint

Version définitive du manuscrit publié dans / Final version of the manuscript published in :  
Plant breeding, 2010, 129 (5), 465-471 <http://dx.doi.org/10.1111/j.1439-0523.2009.01740.x>

- -  
- -

Manuscrit d'auteur / Author Manuscript

Manuscrit d'auteur / Author Manuscript

Manuscrit d'auteur / Author Manuscript

## Figure legend

Fig. 1 Pedigree of the winter wheat line RE714

Fig. 2 Framework genetic maps of groups 2A3, 5D, 6A2 and 7A3 obtained in the wheat RE714 x Hardi RIL population ((Muranty et al. 2009)). Probable positions of other loci are indicated on the right. The position of the QTL detected for powdery mildew resistance are indicated at right, with a horizontal line representing the peak of each QTL and a vertical line representing the confidence interval of each QTL. Markers for which RE714 had the same allele as VPM on linkage group 2A3 are in red, those for which RE714 had the same allele as *Ae. tauschii* n°33 on linkage group 5D are in green, those for which RE714 had the same allele as *T. dicoccum* n°119 on linkage group 6A2 are in blue and those for which RE714 had the same allele as Beauchamp on linkage group 7A3 are in orange. The markers for which RE714 had the same allele as *T. dicoccum* n°119 on linkage group 7A3 are in blue italics.



