



Improvement of in vitro donor plant competence to increase de novo shoot organogenesis in rose genotypes

Latifa Hamama, Linda Voisine, Sandrine Pierre, Denis Cesbron, Laurent L. Ogé, M. Lecerf, S. Cadieux, J Bosselut, Séverine Foucrier, Fabrice Foucher, et al.

► To cite this version:

Latifa Hamama, Linda Voisine, Sandrine Pierre, Denis Cesbron, Laurent L. Ogé, et al.. Improvement of in vitro donor plant competence to increase de novo shoot organogenesis in rose genotypes. *Scientia Horticulturae*, 2019, 252, pp.85-95. 10.1016/j.scienta.2019.03.040 . hal-02278794

HAL Id: hal-02278794

<https://institut-agro-rennes-angers.hal.science/hal-02278794>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Improvement of in vitro donor plant competence to increase de novo shoot organogenesis in rose genotypes

Hamama L, Voisine L., Pierre S., Cesbron D., Ogé L., Lecerf M., Cailleux S., Bosselut J., Foucrier S., Foucher F.,
Berruyer R., Sakr S., Hibrand-Saint Oyant L.*

IRHS, INRA, Agrocampus-Ouest, University of Angers, SFR 4207 QuaSaV, 49071, Beaucouzé, France

Keywords: *Rosa*, shoot regeneration, Carbohydrate source

***Corresponding author**

E-mail address: laurence.hibrand-saint-oyant@inra.fr,

ABSTRACT

A procedure was developed for *in vitro* propagation of *Rosa* genotypes along with an efficient *de novo* shoot organogenesis (DNSO) method. We tested, on one genotype (hybrid of *Rosa wichurana*), the effects of MS basal medium complemented with two growth regulators to achieve either shoot elongation or shoot multiplication of plants. These media were complemented with carbohydrate concentrations from different sources. Then, the impacts of various carbohydrates (fructose, glucose, maltose, sorbitol, sucrose) on the growth and development of several rose genotypes during donor plant subculturing were studied on SMM. The results showed high variability in growth and development between genotypes. Contrary to other members of the *Rosaceae* family, no correlation was found between the shoot size and number when the amount of sorbitol was increased.

Murashige and Skoog medium supplemented with 3.0 mg.L⁻¹ BAP and containing fructose or glucose at 30 g.L⁻¹ was chosen to induce leaf explants for the DNSO experiments. MS basal medium complemented with TDZ/IBA at three ratios and the same range of carbohydrate sources were tested for DNSO. Significant genotypic variations with regard to the percentage of regeneration was demonstrated with six genotypes. For two genotypes, a hybrid of *Rosa wichurana* and *Rosa* 'White Pet', we defined the conditions required to obtain 100% DNSO. For *Rosa chinensis* 'Old blush' and the rootstock genotype *Rosa* 'Natal Briar', we obtained 74 and 87.5% DNSO and only 56.67% and 37.5% for *Rosa* GUY SAVOY® ('Delstrimen') and *Rosa* 'Félicité et Perpétue' respectively. This adventitious shoot regeneration method may be used for large-scale shoot propagation and genetic engineering studies in *Rosa*.

1. Introduction

Rose is an important plant in the ornamental field but also in cosmetics and food industry. Rose is the most important economically species with production of cut flowers, potted roses and garden roses. Thanks to its broad diversity and high quality genome sequence, rose is increasingly seen as a model plant for woody ornamentals (Hibrand-Saint Oyant *et al.*, 2018; Mujib *et al.*, 2013). Major ornamental traits such as scent, flowering and *in vitro* production can be studied in rose.

Rose varieties are conventionally propagated by cuttings or grafting onto a rootstock. The history of *in vitro* rose culture started in 1945 (Nobecourt and Kofler), and since then many reports have documented simple and rapid rose micropropagation methods.

Biotechnology approaches are increasingly used to improve horticultural crop production (Chebet *et al.*, 2003), to boost production by shortening the production time and generating healthy, disease-free plants. *In vitro* multiplication could be used for rapid and mass propagation of cut flowers (Bao *et al.*, 2009) or pot plant production (Fotopoulos and Sotiropoulos, 2004). Martin (1988) demonstrated that up to 400,000 plants could be annually cloned, from a single rose using this technology. Despite the availability of successful micropropagation techniques, *de novo* regeneration methods involving organogenesis (process of forming new organs) and somatic embryogenesis (embryo derived from a somatic cell) remain challenging. Generally, two key pathways can lead to the regeneration of new plants from *in vitro* cultured explants or single cells. Organogenesis and somatic embryogenesis are essential and critical tools for plant multiplication, crop improvement, plant functional genomics and genome editing.

The organogenesis system, also called *de novo* shoot organogenesis (DNSO) (Duclercq *et al.*, 2011), refers to the capacity to regenerate a new tissue culture plant from somatic cells. The advantages of DNSO are a short callus phase, which maintains uniformity, and a reduction in somaclonal variation, often derived from the callus, suspension or protoplasts (for review, see Mujib *et al.* 2013). However, some somaclonal variations in terms of habit and leaf shape have been observed on adventitious shoots from *Rosa persica* x *xanthina* explants (Lloyd *et al.*, 1988).

The prospect of developing a reliable protocol has fostered a tremendous amount of work geared towards identifying the key factors influencing regeneration (more than 3500 papers since 1975 with about 60 on rose). Plant growth regulators (PGR) have been extensively studied, particularly auxin and cytokinin. The aim has not yet been fulfilled, particularly in some so-called 'recalcitrant' species.

In general, plant regeneration in rose is obtained by the embryogenesis pathway rather than organogenesis, although the first report of the regeneration process indicated it was obtained by organogenesis (Hill, 1967). The first report on embryogenesis in rose was published in 1995 by Roberts *et al.* Then several teams developed this regeneration technique, focused on the cultivated species, often tetraploid ones (Dohm *et al.*, 2002; Estabrooks *et al.*, 2007; Kintzios *et al.*, 1999; Xing *et al.*, 2014). In parallel, several studies on organogenesis have been conducted that were focused on the kind of explants, such as leaves or leaflets, roots, internodes and petioles (Arene *et al.*, 1993; Estabrooks *et al.*, 2007; Lloyd *et al.*, 1988; Pati *et al.*, 2004). The organogenesis process in rose was developed either after an induction phase on a cytokinin-rich medium (Dubois *et al.*, 2000; Pati *et al.*, 2004) or on regeneration medium containing cytokinins and auxins (Dubois *et al.*, 2000; Pourhosseini *et al.*, 2013), or with cytokinin alone (Arene *et al.*, 1993; Ibrahim and Debergh, 2001; Lloyd *et al.*, 1988). In all of these studies, regeneration was obtained in the dark or under low light conditions. The roses used in these studies were predominantly cultivated ones, even though some wild species such as *Rosa persica*, *Rosa wichurana*, *Rosa laevigata* (Lloyd *et al.*, 1988), *Rosa chinensis* (Li *et al.*, 2002) and *Rosa damascena* (Pati *et al.*, 2004) have also been used.

Few studies have been conducted to document the strong involvement of sugars in metabolic and developmental processes (for review, see Yaseen *et al.* 2013). Indeed, sugars play a trophic role to sustain the high metabolic activity of heterotrophic organs during growth and act as a signal to control diverse developmental processes (Bolouri Moghaddam and Van den Ende, 2013; Lastdrager *et al.*, 2014; Matsoukas, 2014; Barbier *et al.*, 2015; Xiong *et al.*, 2013).

To further improve the *de novo* shoot organogenesis process in rose genotypes, it is essential to control the physiological state of donor plants in order to prepare explant tissues and increase their regeneration potential. No comparative experiments have been published on the kind of sugars involved in rose mother-plant production or in the regeneration medium.

In vitro plant cells, tissues and organ cultures are mainly heterotrophic structures that are unable to produce their own sugars. Additional exogenous carbohydrates are thus required in the artificial culture media to meet the high energy requirements of developmental processes such as *in vitro* rooting, shoot proliferation and plant regeneration (Barbier *et al.*, 2015; Matsoukas *et al.*, 2013; Yaseen *et al.*, 2013). Many types of carbohydrate, including sucrose, sucrose-derivative hexoses (glucose and fructose) and polyols (sorbitol and mannitol) have been tested in terms of the morphogenesis potential of *in vitro* cultured tissues (for review, see Yaseen *et al.* 2013). Mainly sucrose and sorbitol have been tested as exogenous carbon sources since they are photosynthesis products and phloem-

translocated to different sink organs where they are metabolized (Maurel *et al.* 2004; Gao *et al.* 2003), particularly in the *Rosaceae* family. Yaseen *et al.* (2009) reported that both sucrose and sorbitol play a central role in *in vitro* shoot proliferation in M9 and M26 apple rootstocks. Likewise, sucrose modulates *in vitro* shoot development in cork oak (Romano *et al.*, 1995) and *Eclipta alba* (Baskaran and Jayabalan, 2005). In *Stevia rebaudiana*, shoot proliferation was found to be sensitive to sucrose and fructose (Preethi *et al.*, 2011), while shoot proliferation in *Prunus mume* was more sensitive to glucose (Harada and Murai, 1996). Sucrose is the most widely used carbohydrate source for plant regeneration (Fatima *et al.*, 2015) while mannitol is considered to be a metabolically inert carbohydrate, except in a few species (Conde *et al.*, 2007; Noiraud *et al.*, 2001) and it had very little effect on *in vitro* shoot development and even led to cell death in soybean explants (Sairam *et al.*, 2003). Taken together, these findings indicate that the choice of carbohydrates and/or their concentration could be a powerful lever to successfully manage the morphogenic competence of plant tissue cultures. Furthermore, sugars are increasingly considered as signal entities that are perceived by diverse sensors and regulate many fundamental plant biology processes through a complex regulatory network integrating endogenous (hormones) and exogenous (environment) cues (Broeckx *et al.*, 2016; Moore *et al.*, 2003; Robaglia *et al.*, 2012; Rolland *et al.*, 2006, Sakr *et al.*, 2018). It seems clear that the morphogenic potential of plant tissues could be a complex mechanism and may result from interlinked processes involving sugars, hormones and environmental factors.

The aim of the present research was to develop a strategy to improve the totipotency and organogenic potential, especially the DNSO process, in seven rose genotypes by adjusting the micropropagation method and medium. To determine the specific physiological state that explants should be excised after the clonal cycle in donor plants, we studied the type and amount of carbohydrate source at various regeneration stages, including mother plant production.

2. Materials and Methods

2.1. Plant material and culture conditions

Four diploid cultivars, differing in their growth habit and recurrent blooming capacity, were used in this study. A hybrid of *Rosa wichurana* (Rw) obtained from a rose garden in the “Jardin de Bagatelle” (Paris, France) is a once-flowering genotype with indeterminate vegetative shoots and a ground-cover habit. *Rosa chinensis* ‘Old Blush’ (OB) is a continuous-flowering genotype with a terminal inflorescence and bush habit. The cultivated roses *Rosa* ‘Félicité et Perpétue’ (FP), a once-flowering genotype with indeterminate vegetative shoots and climbing habit, and *Rosa* ‘White Pet’ (WP), historically named ‘Little White Pet’, a sport of FP with a continuous-flowering behaviour (terminal inflorescences and bush habit, Iwata *et al.* 2012). The plants were maintained in a greenhouse in the general following conditions: minimum air temperature maintained at 18°C, with an aeration at 20°C; relative humidity maintained at 70% and no complementary lighting.

The methodology was validated on three other rose genotypes obtained as bare-rooted plants from the “Société Nouvelle des pépinières Georges Delbard” (Malicorne, France). These plants were multiplied in the greenhouse before their *in vitro* introduction.

These varieties are two continuous-flowering genotypes and tetraploid cultivated garden roses ($2x=4n=28$) (*Rosa* PIMPRENELLE® (‘Deldog’) and *Rosa* GUY SAVOY® (‘Delstrimen’)) and one diploid rootstock variety (*Rosa* ‘Natal Briar’).

2.2. Meristems and *in vitro* culture conditions

In order to work with healthy bacteria-free materials, and since *in vitro* mother-plant material is often internally contaminated by bacteria if introduced *in vitro* via node culture (data not shown), meristem cultures were conducted to obtain bacteria-free *in vitro* plants. From the plants maintained in the greenhouse, shoot stems were harvested, surface-sterilized by rapid immersion in 70% ethanol, and meristems (axillary buds) were prepared as described by Lebras *et al.* (2014). Meristems were grown in a growth chamber at $20 \pm 2^\circ\text{C}$ under a 16 h photoperiod (light intensity of $40 \mu\text{mol.m}^{-2}.\text{s}^{-1}$ provided by fluorescent tube lights).

2.3. Clone maintenance and micropropagation

Meristem-derived *in vitro* plants were multiplied and maintained in clonal cycles by subculturing nodes, with each containing two axillary buds. In a first step, explants were cultured on Murashige et Skoog (MS) basal medium complemented with 0.05 g.L^{-1} Fe-EDDHA, 0.1 mg.L^{-1} GA₃ and 0.5 mg.L^{-1} BAP (6-benzylaminopurine),

sucrose 30 g.L⁻¹ and solidified with 3 g.L⁻¹ Phytigel™ (Sigma) (shoot elongation medium, SEM). The medium pH was adjusted to 5.7 before sterilization (113°C, 20 min). The cultures were conducted under a 16 h photoperiod with a photosynthetic flux (PAR) of 56.4 μmol m⁻² s⁻¹ (generated by a combination of two Sylvania Luxline F58W/840 fluorescent lamps and one Osram Biolux L 58W/72-965 fluorescent lamp). The temperature regime was 23 ± 0.5 °C during the 16 h light period and 19 ± 0.5 °C during the 8 h dark period.

Individual Rw shoots were subcultured every month on SEM or SMM (corresponding to an SEM cytokinin enriched medium: 3 mg.L⁻¹ BAP without GA₃), while only SMM was used for the other genotypes. Moreover, on each medium (SEM and SMM), two replicates of a sugar comparison were carried out using various carbohydrate types and concentrations, corresponding to sucrose, glucose, fructose, maltose or sorbitol at 10, 15, 30 or 40 g.L⁻¹. To assess and compare the impact of the medium hormonal and sugar content on plant development, different phenotypic traits were measured after one month on the medium, such as: shoot size (height), shoot number, a scale of the presence or absence of callus (0=absence and 1-2-3-4-5=scale of size callus), the number of new roots, node number and ramification (axillary bud burst).

In vitro plants maintained in a clonal cycle were used as explant sources for the induction of shoot regeneration via organogenesis.

2.4. *De novo* shoot organogenesis

For the Rw genotype, donor plants were subcultured on SEM and SMM, while only SMM was used for the other genotypes.

Explants were excised from 6 week-old *in vitro* plants cultivated on SMM medium with glucose or fructose at 30 g.L⁻¹. The clonal cycle included 1 week in darkness. Explants, consisting of the two proximal leaves and corresponding to the three terminal leaflets with subtending petiolules, were incubated on induction medium and the impact of sugar was compared using 5 types of carbohydrate source (fructose, glucose, maltose, sorbitol and sucrose) at different concentrations (10, 15, 30, 40 g.L⁻¹).

For each condition, two replicates of 10-12 wounded leaf explants were placed with their abaxial side on petri dishes containing solid regeneration MS medium according to Ibrahim and Debergh (2001). Two or three scratches (depending of the leaf maturity and size) were made on the abaxial of each leaflet with a scalpel perpendicular to the midrib. Shoot regeneration medium (SRM) containing macro- and micro-elements, MS vitamins, was complemented with 0.56 mM myo-inositol and solidified with 3 g.L⁻¹ Phytigel. The effects of growth regulator

combinations were evaluated by supplementing the medium with 2.3, 4.6 or 9.2 μM of TDZ (Thidiazuron) (SRM 1, 3 and 6, respectively) and with 0.23 μM (SRM 1 and 3) or 0.46 μM (SRM 6) of IBA using compartmentalized petri dishes, as shown in the Figure 4b. The medium pH was adjusted to 5.7 before sterilization. Cultures were maintained in a tissue culture room under the above described temperature regime.

All cultures were initially incubated in the dark at 23°C for 16 h at 19°C for 8 h for 1 or 2 weeks depending on the time to achieve DNSO.

After 6 weeks, the following explant characteristics were measured: relative size of callus (0 to 5 scale, corresponding to the absence (0) or presence of callus (1 to 5) and depending on the callus diameter (in cm) 1 corresponding to <0.5 ; 2 is 0.5 to 1; 3 is 1 to 1.5; 4 is 1.5 to 2 and 5 is >2), presence or absence of roots, shoot number per explant and shoot size.

The regenerated plants were rooted on MS basal medium supplemented with 0.1 mg. L⁻¹ AIB and 0.5 mg.L⁻¹ NAA at pH 6.

2.5. Statistical analyses

All data were statistically analyzed with the R software package, version 2.11.1 (<http://www.r-project.org/>), including ANOVA, HSD tests for multiple comparisons and logistic analysis. Quantitative data (shoot size, shoot number) were analyzed with ANOVA, followed by a Tukey's HSD post-hoc test. Bimodal DNSO data (i.e. 1 for DNSO and 0 for no DNSO) were fitted in logistic models. 95% confidence intervals were then calculated for each LOD-odds ratio. Log odds ratios were then converted back into DNSO probabilities. The confidence interval was not calculated when no DNSO was observed.

3. Results

To define the best conditions for DNSO on various genotypes, we first tested two mediums and different sugars on the genotype Rw (Fig. 5). The best medium was defined for its potential to induce young tissue. This medium was then used on other genotypes (Fig. 5) and DNSO was conducted on explants which were taken on in vitro plants produced on this best medium.

3.1. Clonal cycle of the hybrid of *Rosa wichurana* (Rw)

A clonal cycle was performed to multiply *in vitro* mother plants and produce explants able to induce *de novo* shoot organogenesis.

In vitro culture of Rw was achieved via meristem culture. The effects of combined growth regulators (SEM and SMM) and carbohydrates (different types and concentrations) on shoot and root development and on callus formation during the clonal cycle were evaluated (illustrated in Table 1 and Fig. 1).

For fructose, glucose and sucrose, we observed an effect-relationship between the increased sugar concentration and the callus size, irrespective of the media (SEM or SMM). Roots developed only on SMM medium with an increased number of shoot correlating with the increase concentration of fructose, glucose and sucrose (Table 1b). In this respect, no significant relationship was noted for maltose and sorbitol (Table 1a).

Node formation (NoNode) and ramifications were only observed on SEM. With fructose, glucose or sucrose, we observed an increase in the node number and branching until 30 g.L⁻¹ of these sugars, followed by a decrease at higher concentration (40 g.L⁻¹). For maltose and sorbitol, the relationship was less clear but a trend revealed a maximum node number with the minimum quantity of these two sugars (i.e. 10 g.L⁻¹) (Table 1a).

Except for sorbitol, the shoot size and shoot number increased with the increased sugar concentration until 30 g.L⁻¹ (Table 1). For the high concentration (40 g.L⁻¹), we noted a negative effect on the shoot number and shoot size, except for maltose and sorbitol.

No statistically significant difference in shoot size was observed for fructose, glucose and sucrose at 30 g.L⁻¹ regardless of the medium, and for sucrose at 40 g.L⁻¹ with SEM (Table 1). On SEM, the highest shoot number (but without any significant difference) was observed regardless of the sugar and concentration variation, except for maltose at 10 g.L⁻¹, sorbitol at 10-15-30 g.L⁻¹ and sucrose at 30 g.L⁻¹. On SMM, the highest shoot number was observed for fructose 15-30 g.L⁻¹, glucose 15-30 g.L⁻¹, maltose 30-40 g.L⁻¹ and sucrose 30 g.L⁻¹ (Table 1 and Table S1).

Thus, the shoot height (Shoot size) was clearly higher on SEM compared to SMM, while the opposite trend was observed for basal multiplication (Noshoot) (Fig. 1 and Table 1), especially for SMM containing fructose as carbohydrate source. Overall, we observed better Rw shoot development on media containing fructose, glucose or sucrose compared to those containing maltose and sorbitol. The least favorable medium for shoot development (shoot number and size) was that containing 3 mg.L⁻¹ of BAP (SMM) with sorbitol or maltose as carbohydrate source (Fig. 2). In this genotype, the best medium for the shoot number was SMM supplemented with fructose at 30 g.L⁻¹, while the best shoot growth was obtained with SEM supplemented with glucose at 30 g.L⁻¹.

As the aim of this study was to obtain young tissue for DNSO and since SMM was the best medium for new young shoot induction (Table 1), this medium was chosen for subsequent analyses.

3.2. Clonal cycle of five rose genotypes

We then opted to evaluate SMM (BAP 3 mg.L⁻¹) on six other genotypes: *Rosa* GUY SAVOY® ('Delstrimen') (GS), *Rosa* PIMPRENELLE® ('Deldog') (P), *Rosa chinensis* 'Old Blush' (OB), *Rosa* 'Félicité et Perpétue' (FP), *Rosa* 'Natal Briar' (NB) and *Rosa* 'White Pet' (WP).

GS appeared to be the most productive genotype in terms of growth (Shoot size of 2,75) and new shoot formation (shoot number = NoShoot) (Table 2) up to 6.4 shoots using fructose at 15 g.L⁻¹. For GS and P, in contrast to Rw, we observed a trend towards a negative impact of the sugar concentration (fructose, sorbitol or sucrose) on the shoot size, i.e. the shoot size decreased as the sugar concentration increased. For FP, there were few significant differences in the shoot size and number relative to the sugar type and concentration. Overall, except for GS, sorbitol, regardless of the concentration and genotype, was less effective in promoting shoot growth and stem multiplication (Table 2). Whatever the genotype, the best outcomes, in terms of multiplication level and *in vitro* growth, were obtained with fructose and glucose at concentrations between 10 and 30 g.L⁻¹ (except WP).

Some preliminary analyses were performed on the DNSO potential of explants from the two media and in various sugar types and concentrations (data not shown). The results showed that no DNSO was obtained in explants from SEM and that sucrose and fructose were the most favorable sugars.

Therefore, for further DNSO experiments, we decided to select sucrose at 30 g.L⁻¹ or fructose at 30 g.L⁻¹ as the carbon source and SMM for explant production.

3.3. De novo shoot organogenesis of *Rosa wichurana*

We used 6 week-old Rw *in vitro* plants (cultivated 5 weeks under light conditions then one week under dark conditions). Explants (leaves) from SMM sucrose at 30 g.L⁻¹ or SMM fructose at 30 g.L⁻¹ were placed on SRM1, 3 or 6. On these three media, the same range of sugars (type and concentration) as those used in the clonal cycle were tested. All regeneration was observed 6 weeks after limited callus formation on tissue, indicating an indirect regeneration process.

As shown in Fig. 4a, we first observed callus induction on the petiole or petiolule and on the leaf scratches. Calli were creamy and light brown in color. We observed the first shoot organogenesis right after this callus induction between 3 to 6 weeks (Fig. 4c).

No differences in DNSO process were observed between the sugar types (sucrose or fructose) (data not shown). The effect of the medium and sugar on the mean DNSO percentage is shown in Fig. 2. No significant differences were observed among the tested media (SRM1-SRM3-SRM6) for this genotype, although SRM6 seemed to display a slightly lower DNSO percentage, showing that the increase in auxin (IBA) and cytokinin (TDZ) could be detrimental to the DNSO process for this genotype, regardless of a comparable growth regulator ratio between SRM3 and SRM6. Despite the rise in cytokinin concentration between SRM1 and SRM3, no significant difference in the DNSO percentage was observed.

The maximum DNSO percentage (100%) was observed using sucrose at 30 g.L⁻¹ on SMM during the clonal cycle, then various SRM and sugars in the DNSO medium (Table 3). Fig. 2b shows that higher DNSO percentages were obtained for fructose (85 to 92%) and glucose (75 to 88%) regardless of the concentration, and for glucose at 30 or 40 g.L⁻¹. The shoot regeneration frequency decreased from a mean of 77% to 40% when maltose was increased up to 4%, while no or very low (9%) regeneration was recorded with sorbitol. Regenerated plants were rooted (Fig 4i-j), acclimated (Fig. 4k-l) and observed in the greenhouse (Fig. 4m). There were no discernable differences between mother and regenerated plants.

3.4. Application to other genotypes

DNSO was improved on the five other corresponding genotypes: GS, OB, FP, NB and WP. All genotypes originated from meristem cultures, were subcultured on fructose or sucrose at 30 g.L⁻¹ in SMM culture for 6 weeks under light/dark conditions, then 1 week under dark conditions (Fig. 3 and Table S3) before DNSO assays. Under our conditions, we observed an indirect DNSO process with a short callus phase for all of these genotypes, as already described for Rw.

For GS, only fructose, glucose and sucrose on SRM media were tested. In all tested conditions (media or sugars), the DNSO percentage was low, with no significant differences (Fig. 3). The highest percentage, whichever the medium, was noted with sucrose at 30 and 40 g.L⁻¹, with a mean of 27% and 26%, respectively, and then 19% for fructose at 30 g.L⁻¹ and glucose at 30 and 40 g.L⁻¹ (Fig. 3). The best conditions for achieving 57% DNSO for this genotype involved producing plants on SMM medium containing sucrose at 30 g.L⁻¹ and used the SRM3 medium containing fructose at 30 g.L⁻¹.

We observed high intra-genotype variability for **OB**, for which the best medium seemed to be SRM3, but the differences were not significant (Fig. 3). The best DNSO levels were observed with fructose at 30 g.L⁻¹ (46%) and 40 g.L⁻¹ (40%). No regeneration was achieved with maltose or sorbitol. The maximum DNSO (74%) was obtained for this genotype using fructose at 30 g.L⁻¹ to produce explants and the SRM3 containing fructose at 40 g.L⁻¹.

The lowest DNSO percentages were obtained with the **FP** genotype, for which the highest DNSO level was 37.5% (Table S3) when using sucrose at 30 g.L⁻¹, then SRM1 with fructose at 30 g.L⁻¹. We also observed high variability for this genotype. No regeneration was obtained with sorbitol (Fig. 3).

For NB, like GS, only fructose, glucose and sucrose were tested. The mean DNSO percentage for any given media ranged from 10.5% for sucrose at 15 g.L⁻¹ to 37% for fructose at 40 g.L⁻¹. We observed a slight increase in the DNSO percentage correlated with the increased sugar concentration (Fig.3). Therefore, the highest DNSO percentages were obtained with fructose (62.5%) and sucrose (55.6%) at 40 g.L⁻¹ in SRM6 and SRM1, respectively (Tables 3 and S2).

WP had a different behaviour than that of the genotype from which it was derived (FP), which had a low DNSO capacity. Indeed, for WP, we observed 100% DNSO with glucose at 15 and 30 g.L⁻¹ on SRM1 and SRM3, respectively, although we observed a decrease in the DNSO percentage on the other tested media (Fig. 3 and Tables 3 and S3).

Taken together, all of these results showed various behaviors that were genotype-dependent and sugar-dependent with regard to their DNSO capacities in the regeneration process.

4. Discussion

The aim of this study was to develop a protocol for rose DNSO with various steps corresponding to the initiation of *in vitro* culture, then *in vitro* development and plant regeneration, as summarized in the Figure 5. For this study, we observed the effects of the sugar concentration on *in vitro* growth, *in vitro* behaviour and the DNSO process in combination with various PGR on six rose genotypes. The genotypes we used, differed according to their habit, blooming mode, ploidy level and origin, or to their use (rootstock versus scion; old versus modern genotypes). None of these differences were found to be correlated with the behaviour observed in the *in vitro* clonal cycle, sugar use or DNSO process.

Between both tested media for plant elongation (SEM) and multiplication (SMM), basal multiplication was observed on the SMM containing a greater amount of BAP and no rooting, contrary to the situation on SEM. The carbon source and concentration were also tested. The carbon source is essential to sustain independent *in vitro* growth and organogenesis as has been shown, particularly in strawberry, a species closely related to rose (Grout and Price, 1987). This improved regenerative behaviour has not always been linked to the carbohydrate nutritional status, but rather to the osmotic condition induced by carbohydrates (Gaj, 2004; Lou and Kako, 1995; Nakagawa *et al.*, 2001). Sucrose is often assumed to be the sugar of choice in cell and tissue culture media because it is the predominant sugar that is translocated in phloem (Brachi *et al.*, 2010; Faure *et al.*, 1998; Marino *et al.*, 1993; Pua and Chong, 1985; Sharma *et al.*, 2008), and also due to its cheap and easy availability. In our study, we showed that sucrose was not always the best carbohydrate and the explants grew well on media supplemented with other carbohydrates, such as fructose and glucose. This is comparable to results obtained in other species, such as *Alnus* spp. (Welander *et al.*, 1989) and *Quercus* (Romano *et al.* 1995), which grow better on medium supplemented with glucose, whereas fructose is superior to sucrose for *in vitro* culture of *Morus alba* (Oka and Ohyama, 1986), *Castanea sativa* (Chauvin and Salesses, 1988) and *Malus* rootstock (Welander *et al.* 1989).

Obviously, whilst sucrose is generally applied in tissue culture of plants, its effects on plant development, photosynthetic performance and growth under these conditions seems to be variable and species dependent. Sugar and starch dynamics in the medium-root-leaf system highlight possibilities for optimizing plant tissue culture. Following invertase activity, sucrose is split into glucose and fructose (Straus, 1962). This hydrolysis can take place both inside (within cells by cytosolic or vacuolar invertases) and outside the plant (in *in vitro* medium via cell wall invertase). The amount of hydrolyzed sucrose and its reaction rate are species dependent (George, 1993). Once hexoses are present in the medium (by adding it as the preferable carbon source or by invertase action on sucrose), these sugars can also be used by *in vitro* plants (George, 1993; Wyse, 1979).

However, our results showed that sorbitol, a sugar alcohol, is detrimental to growth and organogenesis in any rose tested genotypes, contrary to what has been previously shown in several studies on various members of the *Rosaceae* family, in which sorbitol is an optimum source of carbon, energy and osmotic adjustment (Kadota and Niimi, 2004; Marino *et al.*, 1993; Pua and Chong, 1984; Sotiropoulos *et al.*, 2006; Yaseen *et al.*, 2013). In apricot microshoots (Marino *et al.* 2010), there are specific enzymes for sorbitol oxidation, including sorbitol dehydrogenase, which convert sorbitol to fructose, thereby improving shoot production and development when sorbitol is added to the medium. This is not true for *Rosa* spp that is known to be deprived of sorbitol-metabolizing enzymes. In other species, sorbitol acts mainly as an osmotic regulator and, unlike sucrose, sorbitol neither supports *in vitro* shoot growth nor is metabolized in most higher plants (Yaseen *et al.* 2013). On the other hand, sorbitol was reported to be completely ineffective in stimulating shoot proliferation and root induction in *Quercus suber* (Romano *et al.* 1995).

Our results showed that maltose yielded low results compared to fructose, glucose and sucrose. Maltose is derived from starch degradation and is a carbon source while also being an osmotic agent. Such a low effect of maltose could be assigned to its weak intake and/or metabolism by *in vitro* cultured tissue. Few studies have been carried out on the effect of maltose on the micropropagation process, Bahmani *et al.* (2009) showed that the size and shoot number of M9 apple rootstock were positively affected using maltose in *in vitro* culture media.

In this study, we also tested the impact of the concentration of various sugars on *in vitro* growth and development. Source activities (e.g. photosynthesis, export) are upregulated under low sugar concentrations whilst sink processes (e.g. growth, storage) are upregulated when carbon sources are abundantly present (Rolland *et al.*, 2006). Lembrechts *et al.* (2017) concluded that, in horticultural production, higher carbohydrate contents (starch, hexoses and sucrose) have substantial impacts since lower enriched media (e.g. 5 g.L⁻¹ instead of 25 g.L⁻¹) could potentially be used to grow healthy *in vitro* plants able to perform photosynthesis immediately when transferred to the greenhouse, while not compromising plant development and growth during the entire production cycle. Concerning plant growth on medium supplemented with fructose, glucose and sucrose, we observed two trends, either an increase or decrease depending on rose genotype in relation to the sugar concentration in the 10 g.L⁻¹ to 30 g.L⁻¹ range. This indicates that plant growth is a complex phenomenon that not only depends on the nature and concentration of the sugars involved but also on the genotype of interest. Concerning organogenesis, it can be concluded that optimal concentrations for the highest number of new shoots was 30 g.L⁻¹ for all sugars, except sorbitol for two genotypes (GS and NB) for which the shoot number decreased with increasing sorbitol content. In general, and regardless of the sugars involved, the highest tested concentration (40 g.L⁻¹) was detrimental to the

growth and development of all genotypes. Using 'Red Globe' grape plantlets, Mao *et al.* (2018) suggested that lower glucose concentrations (10-20 g.L⁻¹) promoted photosynthesis and growth, and the decrease in the photosynthesis rate at higher concentrations (40 g.L⁻¹) may be due to the downregulation of rubisco activity. In line with this, Capellades *et al.* (1991) found that the difference between the photosynthesis rates of test tube *Rosa* (*Rosa multiflora* L. cv.) seedlings grown at 1 and 3% sucrose concentrations were not significant, whereas they decreased significantly at 5%. This rubisco downregulation in response to the accumulation of soluble sugars in leaves could be mediated by a hexokinase signaling dependent pathway (Dai *et al.*, 1999). High carbon levels may also affect cellular growth by affecting the water potential of the medium, which is a very important factor since it determines the movement of water and mineral elements into plant tissues and also maintains better turgor for the plant cells.

To improve *de novo* shoot organogenesis, we produced explants from plants multiplied on medium containing sugar that ensured a maximum of tested genotypes and a higher shoot number, i.e. glucose and fructose at 30 g.L⁻¹.

Several previous studies on DNSO (direct or indirect) showed that leaves (leaflets or trifoliate structure) and petioles are the best tissues to induce *de novo* shoot organogenesis (Arene *et al.* 1993; Dubois *et al.* 2000; Li *et al.* 2002; Lloyd *et al.* 1988; Pourhosseini *et al.* 2013). Only one study dealt with the positive impact of sugar (glucose versus sucrose) on rose DNSO (Hsia and Korban, 1996) on one genotype and no significant difference for another one. Our results clearly showed high variability in the DNSO potential of the tested genotypes, as shown by Dubois *et al.* (2000) and Nguyen *et al.* (2017).

In the present study, we highlighted optimal conditions to obtain 100% DNSO for three genotypes (RW and WP) and the highest percentage for the four others (NB 87.5%; OB 74%, GS 56.7% and FP 37.5%). The average regeneration rate per genotype ranged from 37.5% to 100%, whereas other studies indicated a range of 62% to 100% on 24 genotypes (Dubois *et al.* 2000) and 0.88% to 88.33% on 96 genotypes (Nguyen *et al.* 2017). The origin of this variability is unknown and may be dependent on the genotype. By a GWAS approach, Nguyen *et al.* (2017) found SNP markers linked to this variability and listed candidate genes for shoot regeneration. The superiority of sucrose and its derivative hexoses (glucose, fructose) in promoting shoot regeneration from leaf explants is consistent with the results obtained with rose cultivars and *R. chinensis minima* (Hsia and Korban 1996), and with *Annona muricata* (Lemos and Baker, 1998). Conversely, for *Solanum aculeatissimum*, the highest regeneration percentage was obtained with 30 g.L⁻¹ sucrose (Ghimire *et al.*, 2012), and similar results have been reported for *Echinacea angustifolia* (Kim *et al.*, 2010) and *Harpagophytum* sp. (Jain

et al., 2009). Fructose and maltose, on the other hand, were found to be better carbon sources than sucrose in *Juglans regia* (Seo et al., 2010). These findings suggest that the nature of the carbon source in the culture medium differs among plants species and is a key factor to successfully manage the organogenic competence of plant tissue cultures. These findings pave the way for new investigations into the exact role of sugars in these developmental aspects of plant tissue culture.

In this study, a reliable regeneration system was developed with rose leaves. This method could be used for rapid propagation and genetic transformation studies of this ornamental species.

Declaration of interest The authors declare that they have no conflict of interest.

Acknowledgements

The authors thank Rémi Gardet and his team of the IRHS experimental station (ImHorPhen). They also thank Tatiana Thouroude for technical assistance in plant management and the students Merick Machouri, Mélanie Guideau-Béreau and Julien Chevalier for *in vitro* culture technical assistance. We acknowledge Noëlle Dorion from Agrocampus Ouest for her contribution, David Manley for proofreading the English in this manuscript. This project is funded by the French *Agence Nationale de la Recherche* (Program ANR-11-BTBR-001-GENIUS).

405 **References**

- 406 Arene, L., Pellegrino, C., Gudin, S., 1993. A comparison of the somaclonal variation level of *Rosa hybrida* L. cv.
407 Meirutral plants regenerated from callus or direct induction from different vegetative and embryonic tissues.
408 Euphytica 71, 83-90.
- 409 Bahmani, R., Karami, O., Gholami, M., 2009. Influence of Carbon Sources and Their Concentrations on Rooting
410 and Hyperhydricity of Apple Rootstock MM.106 World Applied Sciences Journal 6, 1513-1517.
- 411 Bao, Y.H., Dharmawardhana, P., Mockler, T.C., Strauss, S.H., 2009. Genome scale transcriptome analysis of shoot
412 organogenesis in *Populus*. BMC Plant Biology 9:132.
- 413 Barbier, F., Peron, T., Lecerf, M., Perez-Garcia, M.D., Barriere, Q., Rolcik, J., Boutet-Mercey, S., Citerne, S.,
414 Lemoine, R., Porcheron, B., Roman, H., Leduc, N., Le Gourrierc, J., Bertheloot, J., Sakr, S., 2015. Sucrose is an
415 early modulator of the key hormonal mechanisms controlling bud outgrowth in *Rosa hybrida*. Journal of
416 Experimental Botany 66, 2569-2582.
- 417 Baskaran, P., Jayabalan, N., 2005. An efficient micropropagation system for *Eclipta alba*—A valuable medicinal
418 herb. In Vitro Cellular & Developmental Biology - Plant 41, 532-539.
- 419 Bolouri Moghaddam, M.R., Van den Ende, W., 2013. Sugars, the clock and transition to flowering. Frontiers in
420 Plant Science 4:22.
- 421 Brachi, B., Faure, N., Horton, M., Flahauw, E., Vazquez, A., Nordborg, M., Bergelson, J., Cuguen, J., Roux, F.,
422 2010. Linkage and association mapping of *Arabidopsis thaliana* flowering time in nature. PLoS Genetics 6,
423 e1000940.
- 424 Broeckx, T., Hulsmans, S., Rolland, F., 2016. The plant energy sensor: evolutionary conservation and divergence
425 of SnRK1 structure, regulation, and function. Journal of Experimental Botany 67, 6215-6252.
- 426 Capellades, M., Lemeur, R., Debergh, P., 1991. Effects of sucrose on starch accumulation and rate of
427 photosynthesis in *Rosa* cultured *in vitro*. Plant Cell, Tissue and Organ Culture 25, 21-26.
- 428 Chaney, L., Sharp, A.R., Evans, C.R., Udall, J.A., 2016. Genome Mapping in Plant Comparative Genomics.
429 Trends in Plant Science 21, 770-780.
- 430 Chauvin, J.E., Salesses, G., 1988. The effect of fructose on chestnut micropropagation *Castanea* sp. Comptes
431 Rendus De L Academie Des Sciences Serie Iii-Sciences De La Vie-Life Sciences 306, 207-212.
- 432 Chebet, D.K., Okeno, J.A., Mathenge, P., 2003. Biotechnological approaches to improve horticultural crop
433 production in Kenya, In: Hammerschlag, F.A., Saxena, P. (Eds.), Biotechnology in Horticultural Crop
434 Improvement: Achievements, Opportunities and Limitations, pp. 473-477.
- 435 Conde, C., Silva, P., Agasse, A., Lemoine, R., Delrot, S., Tavares, R., Geros, H., 2007. Utilization and transport
436 of mannitol in *Olea europaea* and implications for salt stress tolerance. Plant and Cell Physiology 48, 42-53.
- 437 Dai, N., Schaffer, A., Petreikov, M., Shahak, Y., Giller, Y., Ratner, K., Levine, A., Granot, D., 1999.
438 Overexpression of *Arabidopsis* hexokinase in tomato plants inhibits growth, reduces photosynthesis, and induces
439 rapid senescence. Plant Cell 11, 1253-1266.
- 440 Dohm, A., Ludwig, C., Schilling, D., Debener, T., 2002. Transformation of roses with genes for antifungal proteins
441 to reduce their susceptibility to fungal diseases, Acta Horticulturae, 52,105-111.
- 442 Dubois, L.A.M., de Vries, P.d., Koot, A., 2000. Direct shoot regeneration in the rose: genetic variation of cultivars.
443 Gartenbauwissenschaft 65, 45-49.
- 444 Duclercq, J., Sangwan-Norreel, B., Catterou, M., Sangwan, R.S., 2011. De novo shoot organogenesis: from art to
445 science. Trends in Plant Science 16, 597-606.
- 446 Estabrooks, T., Browne, R., Z.M, D., 2007. 2,4,5-Trichlorophenoxyacetic acid promotes somatic embryogenesis
447 in the rose cultivar 'Livin' Easy' (*Rosa* sp.). Plant Cell Reports 26, 153-160.
- 448 Fatima, N., Ahmad, N., Ahmad, I., Anis, M., 2015. Interactive Effects of Growth Regulators, Carbon Sources, pH
449 on Plant Regeneration and Assessment of Genetic Fidelity Using Single Primer Amplification Reaction (SPARS)
450 Techniques in *Withania somnifera* L. Applied Biochemistry and Biotechnology 177, 118-136.
- 451 Faure, O., Diemer, F., Moja, S., Jullien, F., 1998. Mannitol and thidiazuron improve *in vitro* shoot regeneration
452 from spearmint and peppermint leaf disks. Plant Cell Tissue and Organ Culture 52, 209-212.
- 453 Fotopoulos, S., Sotiropoulos, T.E., 2004. In vitro Propagation of the Peach Rootstock: The Effect of Different
454 Carbon Sources and Types of Sealing Material on Rooting. Biologia Plantarum 48, 629-631.
- 455 Gaj, M.D., 2004. Factors influencing somatic embryogenesis induction and plant regeneration with particular
456 reference to *Arabidopsis thaliana* (L.) Heynh. Plant Growth Regulation 43, 27-47.
- 457 Gao, Z.F., Maurousset, L., Lemoine, R., Yoo, S.D., van Nocker, S., Loescher, W., 2003. Cloning, expression, and
458 characterization of sorbitol transporters from developing sour cherry fruit and leaf sink tissues. Plant Physiology
459 131, 1566-1575.
- 460 George, E.F., 1993. Plant propagation by tissue culture. Exegetics Ltd.
- 461 Ghimire, B.K., Yu, C.Y., Chung, I.M., 2012. Direct shoot organogenesis and assessment of genetic stability in
462 regenerants of *Solanum aculeatissimum* Jacq. Plant Cell Tissue and Organ Culture 108, 455-464.

Grout, B.W.W., Price, F., 1987. The establishment of photosynthetic independence in strawberry cultures prior to transplanting., Proc. Symposium Florizel 87, Arlon, Belgium, pp. 55-60.

Harada, H., Murai, Y., 1996. Micropropagation of *Prunus mume*. Plant Cell Tissue and Organ Culture 46, 265-267.

Hill, G.P., 1967. Morphogenesis of shoot primordia in cultured stem tissue of a garden rose. Nature 216, 596-597.

Hsia, C.N., Korban, S.S., 1996. Organogenesis and somatic embryogenesis in callus cultures of *Rosa hybrida* and *Rosa chinensis minima*. Plant Cell Tissue and Organ Culture 44, 1-6.

Ibrahim, R., Debergh, P.C., 2001. Factors controlling high efficiency adventitious bud formation and plant regeneration from in vitro leaf explants of roses (*Rosa hybrida* L.). Scientia Horticulturae 88, 41-57.

Iwata, H., Gaston, A., Remay, A., Thouroude, T., Jeauffre, J., Kawamura, K., Hibrand-Saint Oyant, L., Araki, T., Denoyes, B., Foucher, F., 2012. The *TFL1* homologue *KSN* is a regulator of continuous flowering in rose and strawberry. Plant Journal 69, 116-125.

Jain, N., Bairu, M.W., Stirk, W.A., Van Staden, J., 2009. The effect of medium, carbon source and explant on regeneration and control of shoot-tip necrosis in *Harpagophytum procumbens*. South African Journal of Botany 75, 117-121.

Kadota, M., Niimi, Y., 2004. Influences of carbon sources and their concentrations on shoot proliferation and rooting of 'Hosui' Japanese pear. Hortscience 39, 1681-1683.

Kim, J.S., Lee, S.Y., Eom, S.H., Park, S.U., 2010. Improved shoot organogenesis and plant regeneration of *Echinacea angustifolia* DC. Journal of Medicinal Plants Research 4, 587-591.

Kintzios, S., Manos, C., Makri, O., 1999. Somatic embryogenesis from mature leaves of rose (*Rosa* sp.). Plant Cell Reports 18, 467-472.

Lastdrager, J., Hanson, J., Smeeckens, S., 2014. Sugar signals and the control of plant growth and development. Journal of Experimental Botany 65, 799-807.

Le Bras, C., Le Besnerais, P.H., Hamama, L., Grapin, A., 2014. Cryopreservation of ex-vitro-grown *Rosa chinensis* 'Old Blush' buds using droplet-vitrification and encapsulation-dehydration. Plant Cell Tissue and Organ Culture 116, 235-242.

Lembrechts, R., Ceusters, N., De Proft, M.P., Ceusters, J., 2017. Sugar and starch dynamics in the medium-root-leaf system indicate possibilities to optimize plant tissue culture. Scientia Horticulturae 224, 226-231.

Lemos, E.E.P., Baker, D.A., 1998. Shoot regeneration in response to carbon source on internodal explants of *Annona muricata* L. Plant Growth Regulation 25, 105-112.

Li, X.Q., Krasnyanski, S.F., Korban, S.S., 2002. Somatic embryogenesis, secondary somatic embryogenesis, and shoot organogenesis in *Rosa*. Journal of Plant Physiology 159, 313-319.

Lloyd, D., Roberts, A.V., Short, K.C., 1988. The induction in vitro of adventitious shoots in *Rosa*. Euphytica 37, 31-36.

Lou, H., Kako, S., 1995. Role of high sugar concentrations in inducing somatic embryogenesis from cucumber cotyledons. Scientia Horticulturae 64, 11-20.

Mao, J., Li, W.F., Mi, B.Q., Ma, Z.H., Dawuda, M.M., Zuo, C.W., Zhang, Y.M., Jiang, X.F., Chen, B.H., 2018. Transcriptome analysis revealed glucose application affects plant hormone signal transduction pathway in "Red Globe" grape plantlets. Plant Growth Regulation 84, 45-56.

Marino, G., Bertazza, G., Magnanini, E., Altan, A.D., 1993. Comparative effects of sorbitol and sucrose as main carbon energy-sources in micropropagation of apricot. Plant Cell Tissue and Organ Culture 34, 235-244.

Martin, C., 1985. Plant breeding in vitro, Endeavour ed. 9, 81-86.

Matsoukas, I.G., 2014. Interplay between sugar and hormone signaling pathways modulate floral signal transduction. Frontiers in Genetics 5, 218.

Matsoukas, I.G., Massiah, A.J., Thomas, B., 2013. Starch metabolism and antiflorigenic signals modulate the juvenile-to-adult phase transition in *Arabidopsis*. Plant Cell and Environment 36, 1802-1811.

Maurel, K., Leite, G.B., Bonhomme, M., Guilliot, A., Rageau, R., Petel, G., Sakr, S., 2004. Trophic control of bud break in peach (*Prunus persica*) trees: a possible role of hexoses. Tree Physiology 24, 579-588.

Moore, B., Zhou, L., Rolland, F., Hall, Q., Cheng, W.-H., Liu, Y.-X., Hwang, I., Jones, T., Sheen, J., 2003. Role of the Arabidopsis Glucose Sensor HXK1 in Nutrient, Light, and Hormonal Signaling. Science 300, 332-336.

Mujib, A., Banerjee, S., Ghosh, P.D., 2013. Tissue culture induced variability in some horticultural important ornamentals, Biotechnology & Biotechnological Equipment, pp. 213-224.

Nakagawa, H., Saijyo, T., Yamauchi, N., Shigyo, M., Kako, S., Ito, A., 2001. Effects of sugars and abscisic acid on somatic embryogenesis from melon (*Cucumis melo* L.) expanded cotyledon. Scientia Horticulturae 90, 85-92.

Nguyen, T.H.N., Schulz, D., Winkelmann, T., Debener, T., 2017. Genetic dissection of adventitious shoot regeneration in roses by employing genome-wide association studies. Plant Cell Reports 36, 1493-1505.

Nobecourt, P., Kofler, L., 1945. Bouturage de bourgeons de rosier en milieu nutritif aseptique. Bull. Soc. Bot. 92.

Noiraud, N., Maurousset, L., Lemoine, R., 2001. Identification of a mannitol transporter, AgMaT1, in celery phloem. Plant Cell 13, 695-705.

Oka, S., Ohyama, K., 1986. Mulberry (*Morus alba* L.), In: YP, B. (Ed.), Biotechnology in Agriculture and Forestry Trees I, Springer, New York, , pp. 384–392.

Pati, P.K., Sharma, M., Sood, A., Ahuja, P.S., 2004. Direct shoot regeneration from leaf explants of *Rosa damascena* Mill. In Vitro Cellular & Developmental Biology-Plant 40, 192-195.

Pourhosseini, L., Kermani, M.J., Habashi, A.A., Khalighi, A., 2013. Efficiency of direct and indirect shoot organogenesis in different genotypes of *Rosa hybrida*. Plant Cell, Tissue and Organ Culture 112, 101-108.

Preethi, D., Sridhar, T.M., Naidu, C.V., 2011. Direct Shoot Organogenesis from Leaf Explants of *Stevia rebaudiana*. Journal of phytology 3(5), 69-73.

Pua, E.C., Chong, C., 1984. Requirement for sorbitol (d-glucitol) as carbon source for invitro-propagation of *Malus robusta* no-5. Canadian Journal of Botany-Revue Canadienne De Botanique 62, 1545-1549.

Pua, E.C., Chong, C., 1985. Regulation of *in vitro* shoot and root regeneration in macspur apple by sorbitol (d-glucitol) and related carbon-sources. Journal of the American Society for Horticultural Science 110, 705-709.

Robaglia, C., Thomas, M., Meyer, C., 2012. Sensing nutrient and energy status by SnRK1 and TOR kinases. Current Opinion in Plant Biology 15, 301-307.

Roberts, A.V., Yokoya, K., Walker, S., Mottley, J., 1995. Somatic embryogenesis in *Rosa* spp. In: Somatic Embryogenesis in Woody Plants: Vol 2 — Angiosperms. Dordrecht : Springer Netherlands. pp. 277-289.

Rolland, F., Baena-Gonzalez, E., Sheen, J., 2006. Sugar sensing and signaling in plants: Conserved and novel mechanisms, Annual Review of Plant Biology, pp. 675-709.

Romano, A., Noronha, C., Martinsloucao, M.A., 1995. Role of carbohydrates in micropropagation of cork oak. Plant Cell Tissue and Organ Culture 40, 159-167. Physiology and Molecular Biology of Plants 20, 235-240

Sairam, R.V., Franklin, G., Hassel, R., Smith, B., Meeker, K., Kashikar, N., Parani, M., Al Abed, D., Ismail, S., Berry, K., Goldman, S.L., 2003. A study on the effect of genotypes, plant growth regulators and sugars in promoting plant regeneration via organogenesis from soybean cotyledonary nodal callus. Plant Cell Tissue and Organ Culture 75, 79-85.

Sakr, S., Wang, M., Dedaldechamp, F., Perez-Garcia, M. D., Oge, L., Hamama, L., & Atanassova, R. 2018. The Sugar-Signaling Hub: Overview of Regulators and Interaction with the Hormonal and Metabolic Network. International Journal of Molecular Sciences, 19(9) doi: 10.3390/ijms19092506.

Seo, M.S., Takahara, M., Takamizo, T., 2010. Optimization of culture conditions for plant regeneration of *Panicum* spp. through somatic embryogenesis. . Grassland Sci 56, 6–12.

Sharma, K.D., Rathour, R., Sharma, R., Goel, S., Sharma, T.R., Singh, B.M., 2008. In vitro cormlet development in *Crocus sativus*. Biologia Plantarum 52, 709-712.

Sotiropoulos, T.E., Molassiotis, A.N., Mouhtaridou, G.I., Papadakis, I., Dimassi, K.N., Therios, I.N., Diamantidis, G., 2006. Sucrose and sorbitol effects on shoot growth and proliferation in vitro, nutritional status and peroxidase and catalase isoenzymes of M 9 and MM 106 apple (*Malus domestica* Borkh.) rootstocks. European Journal of Horticultural Science 71, 114-119.

Straus, J., 1962. Invertase in cell walls of plant tissue cultures. Plant Physiol 37, 342-348.

Welander, M., Welander, N.T., Brackman, A.S., 1989. Regulation of invitro shoot multiplication in *Syringa, alnus* and *Malus* by different carbon-sources. Journal of Horticultural Science 64, 361-366.

Wyse, R., 1979. Sucrose Uptake by Sugar Beet Tap Root Tissue. Plant Physiology 64, 837-841.

Xing, W., Bao, Y., Luo, P., Bao, M., Ning, G., 2014. An efficient system to produce transgenic plants via cyclic leave-originated secondary somatic embryogenesis in *Rosa rugosa*. Acta Physiologiae Plantarum 36, 2013-2023.

Xiong, Y., McCormack, M., Li, L., Hall, Q., Xiang, C., Sheen, J., 2013. Glucose–TOR signalling reprograms the transcriptome and activates meristems. Nature 496, 181.

Yaseen, M., Ahmad, T., Abbasi, N. A., Hafiz, I. A. 2009. Assessment of apple rootstocks M 9 and M 26 for in vitro rooting potential using different carbon sources. Pakistan Journal of Botany, 412, 769-781.

Yaseen, M., Ahmad, T., Sablok, G., Standardi, A., Hafiz, I.A., 2013. Review: role of carbon sources for in vitro plant growth and development. Molecular Biology Reports 40, 2837-2849.

Fig. 1 Morphological responses of Rw genotype *in vitro* plants after 6 weeks on SEM and SMM containing various sugars.

Fruc: fructose; Gluc: glucose; Malt: maltose; Sorb: sorbitol; Suc: sucrose at various concentrations (10-15-30 or 40 g.L⁻¹).





Fig. 2 Percentage of mean DNSO obtained from leaf explants of the *Rosa wichurana* hybrid excised from plants cultivated on SMM. The graph shows the percentage of mean DNSO on the different shoot regeneration media, i.e. SRM1, SRM3 or SRM6 (**a**), regardless of the sugar type and concentration and the percentage of mean DNSO according to the sugar type and concentration (**b**) and regardless of the medium.

Fruc=fructose, *gluc*=glucose, *malt*=maltose, *sorb*=sorbitol, *suc*=sucrose

15-30-40 indicate the sugar concentration corresponding to 15 g.L⁻¹, 30 g.L⁻¹ and 40 g.L⁻¹, respectively.

a



b









Fig. 3 Percentage ($\pm$ confidence interval) of the mean DNSO obtained from explants of five genotypes excised from plants cultivated on SMM. The graph shows the percentage of mean DNSO on the various media, i.e. SRM1, SRM3 or SRM6 (a) to the sugar type and concentration and the percentage of mean DNSO according to the sugar type and concentration, regardless of the medium (b).

The genotypes are Rosa GUY SAVOY® ‘Delstrimen’ (GS), Rosa ‘Félicité et Perpétue’ (FP), Rosa ‘Natal Briar’ (NB), Rosa chinensis ‘Old Blush’ (OB) and Rosa ‘White Pet’.

Fruc=fructose, Gluc=glucose, Malt=maltose, Sorb=sorbitol, Suc=sucrose

15-30- 40 indicate the sugar concentration, corresponding to 15 g.L-1, 30 g.L-1 and 40 g.L-1, respectively








Fig. 4 Regeneration of various *Rosa* genotypes. **a)** callus formation on *R.wichurana* (Rw) leaf explants derived from donor plants cultivated on SEM during the clonal cycle **b)** compartmentalized petri dishes used to test three SRM. **c-h)** DNSO on various genotypes. **i-j)** Rooting of Rw plants. **k-l)** Acclimatization of Rw and NB plants in Glass Jar. **m)** Regenerated Rw plant in the greenhouse.

GS: *Rosa* GUY SAVOY® ('*Delstrimen*'), WP: *Rosa chinensis* 'Old Blush', *Rosa* 'Félicité et Perpétue', *Rosa* 'Natal Briar' and *Rosa* 'White Pet'




Fig. 5 Schematic diagram of different steps of rose tissue culture to obtain DNSO, different media components and criteria analysis.
The medium (SEM and SMM) impact was evaluated on the genotype *Rosa wichurana* (*Rw*) and the best medium for DNSO was then used for all genotypes.

Table 1 Impact of SEM [BAP 0.5 mg.L⁻¹/GA3 0.1 mg.L⁻¹] (a) and SMM [BAP 3 mg.L⁻¹] (b) combined with various carbohydrate source types and concentrations on the callus formation, growth and development of *in vitro* *Rosa wichurana* (Rw) explants during the clonal cycle. The mean and standard deviation (SD) were calculated from 12 repetitions in two experiments for the height (cm) of shoots (Shoot size), the number of newly formed shoots derived from the base of the original axillary node (NoShoot), the scale of the size of a callus formed (Callus), the number of roots (Root), the number of formed nodes on the elongated stems (NoNode) and the number of axillary bud bursts on elongated stems (Ramification). Means of Shoot size and NoShoot that are not connected by the same letter are significantly different at 0.05 probability level.

a

SEM	Shoot size		NoShoot		Callus		Root		NoNode		Ramification	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
fructose10	1.90 ^{efghi}	0.55	3.63 ^{ab}	1.07	1.37	0.68	0.05	0.23	1.26	1.05	1.33	0.78
fructose15	2.54 ^{cdef}	0.77	3.40 ^{ab}	1.50	1.65	0.92	0.10	0.24	0.95	1.82	2.08	0.85
fructose30	3.43 ^{ab}	1.20	2.55 ^{abcde}	1.48	2.20	1.47	0.95	1.23	2.65	2.96	1.67	0.93
fructose40	2.75 ^{bcde}	0.79	3.35 ^{abc}	1.33	2.45	1.24	1.50	1.81	1.55	1.89	1.75	0.88
glucose10	1.48 ^{ghi}	0.32	3.33 ^{abc}	1.71	0.88	0.54	0.00	0.00	1.46	1.44	2.00	0.00
glucose15	2.20 ^{defg}	0.68	3.29 ^{abc}	1.82	1.14	0.73	0.05	0.22	0.86	1.42	1.80	0.63
glucose30	3.77 ^a	1.57	2.92 ^{abcd}	1.58	1.92	1.31	1.38	1.56	4.67	5.01	2.42	0.76
glucose40	2.60 ^{bcde}	1.13	3.16 ^{abc}	1.89	2.96	1.22	2.08	1.98	1.68	2.29	2.00	0.74
maltose10	1.25 ^{ghi}	0.53	2.00 ^{bcde}	1.03	0.11	0.32	0.00	0.00	1.89	1.13	0.82	0.40
maltose15	1.42 ^{ghi}	0.37	2.40 ^{abcde}	1.19	0.30	0.47	0.00	0.00	0.70	1.08	1.83	0.39
maltose30	1.59 ^{ghi}	0.38	3.42 ^{ab}	2.09	0.30	0.47	0.05	0.22	0.30	0.92	1.83	0.94
maltose40	1.65 ^{fghi}	0.25	3.85 ^a	1.79	0.35	0.49	0.00	0.00	0.85	1.18	1.77	0.93
sorbitol10	1.27 ^{hi}	0.78	1.53 ^{de}	0.70	0.00	0.00	0.00	0.00	2.42	1.30	0.73	0.47
sorbitol15	1.34 ^{ghi}	0.44	1.32 ^e	0.48	0.37	0.50	0.00	0.00	1.32	1.70	0.45	0.52
sorbitol30	1.20 ^{hi}	0.34	1.75 ^{cde}	0.79	0.30	0.47	0.00	0.00	0.85	1.50	0.83	0.72
sorbitol40	1.13 ⁱ	0.23	2.25 ^{abcde}	1.29	0.30	0.47	0.00	0.00	1.55	1.15	1.00	0.00
sucrose10	1.39 ^{ghi}	0.46	2.79 ^{abcde}	1.50	0.92	0.58	0.00	0.00	0.75	0.74	1.33	0.65
sucrose15	2.02 ^{efgh}	0.76	2.75 ^{abcde}	1.42	1.04	0.91	0.00	0.00	0.67	1.24	1.83	0.83
sucrose30	3.02 ^{abcd}	0.93	1.71 ^{de}	1.23	2.08	1.23	0.54	0.86	4.00	4.44	1.17	1.12
sucrose40	3.35 ^{abc}	1.01	2.67 ^{abcde}	1.49	3.00	1.44	1.29	1.33	3.21	3.66	1.33	0.65

b

SMM	Shoot size		NoShoot		Callus		Root		NoNode		Ramification	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
fructose10	1.10 ^{cde}	0.31	3.38 ^{bcdef}	1.47	0.83	0.82	0.00	0.00	0.00	0.00	0.00	0.00
fructose15	1.37 ^{bcd}	0.34	4.46 ^{ab}	1.38	1.21	1.14	0.00	0.00	0.00	0.00	0.00	0.00
fructose30	2.43 ^a	0.63	4.95 ^a	2.13	2.96	1.46	0.00	0.00	0.00	0.00	0.00	0.00
fructose40	1.65 ^b	0.59	3.00 ^{bcdefgh}	1.35	2.96	1.57	0.04	0.20	0.00	0.00	0.00	0.00
glucose10	1.07 ^{cde}	0.24	3.29 ^{bcdefg}	1.37	0.71	0.91	0.00	0.00	0.00	0.00	0.00	0.00
glucose15	1.33 ^{bcd}	0.56	3.75 ^{abcd}	1.15	0.96	0.95	0.00	0.00	0.00	0.00	0.00	0.00
glucose30	2.20 ^a	0.72	3.77 ^{abcd}	1.69	2.63	1.56	0.00	0.00	0.00	0.00	0.00	0.00
glucose40	1.42 ^{bc}	0.59	1.75 ^h	1.22	3.50	1.56	0.00	0.00	0.00	0.00	0.00	0.00
maltose10	1.02 ^{cde}	0.22	2.21 ^{efgh}	0.88	0.04	0.20	0.00	0.00	0.00	0.00	0.00	0.00
maltose15	1.16 ^{cde}	0.32	2.50 ^{cdefgh}	1.29	0.17	0.38	0.00	0.00	0.00	0.00	0.00	0.00
maltose30	1.31 ^{bcde}	0.44	4.10 ^{abc}	1.87	0.08	0.28	0.00	0.00	0.00	0.00	0.00	0.00
maltose40	1.39 ^{bcd}	0.52	4.13 ^{abc}	2.47	0.38	0.50	0.00	0.00	0.00	0.00	0.00	0.00
sorbitol10	0.81 ^e	0.21	1.86 ^{fgh}	1.04	0.14	0.35	0.00	0.00	0.00	0.00	0.00	0.00
sorbitol15	0.88 ^{de}	0.30	1.81 ^{gh}	0.93	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
sorbitol30	0.84 ^{de}	0.18	2.32 ^{defgh}	1.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
sorbitol40	0.79 ^e	0.21	2.00 ^{efgh}	0.93	0.06	0.25	0.00	0.00	0.00	0.00	0.00	0.00
sucrose10	1.07 ^{cde}	0.46	2.33 ^{defgh}	1.24	0.50	0.66	0.00	0.00	0.00	0.00	0.00	0.00
sucrose15	1.28 ^{bcde}	0.43	3.08 ^{bcdefgh}	1.10	0.92	1.06	0.00	0.00	0.00	0.00	0.00	0.00
sucrose30	2.25 ^a	0.75	3.63 ^{abcde}	1.56	2.42	1.38	0.00	0.00	0.00	0.00	0.00	0.00
sucrose40	1.37 ^{bcd}	0.62	2.63 ^{cdefgh}	1.66	2.50	1.72	0.00	0.00	0.00	0.00	0.00	0.00

Table 2 Results of HSD tests on the shoot size and number (NoShoot) relative to the sugar type and concentration for six genotypes (*Rosa* GUY SAVOY® '**Delstrimen**', *Rosa* PIMPRENELLE® 'Deldog', *Rosa chinensis* 'Old Blush', *Rosa* 'Félicité et Perpétue', *Rosa* 'Natal Briar' and *Rosa* 'White Pet' during the clonal cycle on SMM. Means of Shoot size and NoShoot that are not connected by the same letter (colum Groups) are significantly different at 0.05 probability level.

Rosa GUY SAVOY® ('Delstrimen')

Shoot size		
Groups	Treatments	means
a	fructose10	2.75
ab	glucose10	2.361
ab	fructose15	2.308
abc	sorbitol10	2.264
abc	glucose40	2.258
abc	glucose30	2.233
abc	sucrose10	2.227
abc	glucose15	2.204
abc	sucrose15	2.112
abcd	maltose15	2.104
abcd	maltose10	2.077
bcd	sorbitol15	1.9
bcd	fructose30	1.838
bcd	maltose30	1.821
bcd	sucrose40	1.762
bcde	sucrose30	1.638
bcde	fructose40	1.621
cde	maltose40	1.557
de	sorbitol30	1.364
e	sorbitol40	0.9375

Rosa PIMPRENELLE® ('Deldog')

Shoot size		
Groups	Treatments	means
a	glucose15	1.912
a	fructose10	1.86
ab	glucose30	1.786
ab	fructose15	1.75
ab	sucrose10	1.73
ab	fructose30	1.714
ab	maltose30	1.643
ab	glucose10	1.63
ab	glucose40	1.629
ab	sucrose40	1.571
ab	sucrose15	1.475
ab	maltose15	1.457
ab	sorbitol10	1.412
ab	sucrose30	1.357
ab	maltose10	1.35
ab	fructose40	1.2
ab	sorbitol15	1.071
ab	maltose40	1.014
b	sorbitol30	0.9
b	sorbitol40	0.9

Rosa chinensis 'Old Blush'

Shoot size		
Groups	Treatments	means
a	sucrose30	1.633
ab	fructose30	1.442
abc	fructose40	1.333
abc	sucrose10	1.292
abc	glucose15	1.25
abc	glucose30	1.208
abc	fructose10	1.192
abc	maltose30	1.175
abc	fructose15	1.142
abc	sucrose40	1.125
abc	maltose15	1.05
bc	glucose10	0.9917
bc	maltose10	0.9917
bc	glucose40	0.9833
bc	sorbitol30	0.9417
bc	sucrose15	0.875
bc	maltose40	0.8667
bc	sorbitol10	0.8333
bc	sorbitol40	0.8167
c	sorbitol15	0.8

Rosa 'Félicité et Perpétue'

Shoot size		
Groups	Treatments	means
a	glucose40	1.667
a	fructose30	1.533
ab	glucose30	1.367
ab	fructose40	1.356
ab	maltose40	1.273
ab	glucose15	1.1
ab	fructose15	1.083
ab	maltose30	1.027
ab	sucrose30	1
ab	sucrose10	0.9889
ab	sorbitol30	0.9556
ab	maltose15	0.8545
ab	sorbitol15	0.8125
b	sorbitol40	0.6545
b	sucrose40	0.6333
b	sucrose15	0.4714

Rosa 'Natal Briar'

Shoot size		
Groups	Treatments	means
a	glucose30	1.456
ab	fructose10	1.259
ab	glucose15	1.163
abc	fructose15	1.116
abc	glucose10	1.082
abc	fructose30	1.039
abc	fructose40	1.039
abc	maltose10	1.035
abc	maltose15	1.028
abc	maltose30	1.022
abc	sucrose30	0.9611
abc	sorbitol10	0.9412
abc	glucose40	0.9235
abc	sucrose40	0.9
abc	maltose40	0.8778
abc	UK40	0.8667
abc	sorbitol15	0.8158
bc	sucrose15	0.7895
bc	sorbitol30	0.7278
bc	sucrose10	0.7
bc	UK30	0.6
bc	UK15	0.5
c	sorbitol40	0.4944
c	UK10	0.3333

Rosa 'White pet'

Shoot size		
Groups	Treatments	means
a	glucose40	1.892
ab	fructose40	1.627
ab	glucose30	1.673
abc	fructose30	1.28
abc	maltose30	1.344
abc	maltose40	1.42
bc	sucrose40	0.9182
c	sorbitol30	0.7556
c	sucrose30	0.675

NoShoot		
Groups	Treatments	means
a	fructose15	6.423
ab	fructose10	5.545
abc	glucose30	5.208
abcd	fructose30	5.125
abcde	maltose10	5
abcde	sorbitol10	4.818
abcde	glucose15	4.8
abcde	sorbitol15	4.538
bcde	maltose30	4.417
bcde	glucose10	4.217
bcde	maltose15	3.917
bcde	sucrose30	3.875
bcde	sucrose15	3.692
bcdef	sucrose10	3.591
bcdef	sorbitol30	3.5
cdef	glucose40	3.375
def	maltose40	3.174
ef	fructose40	3.125
ef	sucrose40	3.042
f	sorbitol40	1.708

NoShoot		
Groups	Treatments	means
a	sucrose30	2.714
ab	glucose30	2.571
abc	fructose30	2.286
abc	fructose40	2.286
abc	fructose15	2.25
abc	maltose30	2.143
abc	sucrose40	1.857
abc	glucose40	1.714
abc	maltose40	1.571
abc	glucose10	1.5
bc	fructose10	1.4
bc	glucose15	1.375
bc	sucrose15	1.375
bc	maltose15	1.286
c	sorbitol10	1.25
c	sorbitol30	1.143
c	sorbitol40	1.143
c	maltose10	1.125
c	sucrose10	1.1
c	sorbitol15	1

NoShoot		
Groups	Treatments	means
a	fructose30	4.167
ab	sucrose30	3.5
abc	glucose30	3.167
abcd	maltose30	3.083
abcde	glucose40	2.917
abcde	sucrose40	2.917
bcdef	fructose40	2.5
bcdef	maltose40	2.5
cdef	sorbitol40	1.75
def	glucose10	1.583
def	glucose15	1.583
def	maltose15	1.583
ef	fructose15	1.417
ef	sorbitol30	1.417
f	sorbitol15	1.25
f	fructose10	1.167
f	maltose10	1.167
f	sucrose10	1.083
f	sorbitol10	1.001
f	sucrose15	1.001

NoShoot		
Groups	Treatments	means
a	glucose30	3.083
a	fructose15	3
a	fructose30	2.917
a	maltose30	2.364
a	glucose15	2.167
a	glucose40	2.167
a	fructose40	2.111
a	sorbitol15	2
a	sucrose10	2
a	sucrose40	2
a	maltose40	1.818
a	sucrose30	1.571
a	sorbitol30	1.556
a	maltose15	1.545
a	sorbitol40	1.455
a	sucrose15	1.143

NoShoot		
Groups	Treatments	means
a	fructose30	7.778
ab	glucose15	6.684
abc	fructose10	5.529
abcd	glucose30	5.167
abcd	maltose40	5.167
bcd	glucose10	5
bcde	fructose15	4.842
bcde	fructose40	4.778
bcde	maltose30	4.389
bcdef	maltose10	4.176
bcdef	maltose15	4.167
bcdef	sucrose40	4.167
cdef	sucrose30	3.833
cdef	sucrose15	3.737
cdef	glucose40	3.471
cdef	sorbitol10	3.412
cdef	sucrose10	3.353
cdef	UK40	3.333
def	sorbitol15	2.684
def	UK15	2.667
def	UK10	2.333
def	UK30	2.333
ef	sorbitol30	2.222
f	sorbitol40	1.444

Noshoot		
Groups	Treatments	means
a	maltose30	4.222
ab	glucose40	4
abc	fructose40	3.091
bcd	fructose30	2.5
cd	glucose30	2
cd	maltose40	1.9
cd	sorbitol30	2.111
cd	sucrose30	2.125
d	sucrose40	1.455

Table 3 Optimal conditions to obtain the best percentage of DNSO for each tested genotype.
Suc1 corresponds to the sugar type and concentration during the clonal cycle (Suc: sucrose, fruc: fructose; 30: 30 g.L⁻¹)
Medium corresponds to the medium for DNSO
Suc2 corresponds to the sugar type and concentration in DNSO medium
(Fruc: fructose, Gluc: glucose, Suc: sucrose; 15: 15 g.L⁻¹, 30: 30 g.L⁻¹, 40: 40 g.L⁻¹)
Rw: Rosa wichurana, GS: Rosa GUY SAVOY® ('Delstrimen'), OB: Rosa chinensis 'Old Blush', FP: Rosa 'Félicité et Perpétue', NB: Rosa 'Natal Briar', WP: Rosa 'White Pet'.

	Suc1	Medium	Suc2	% DNSO
		SRM1	Fruc15, Gluc30, Suc15	100
Rw	Suc30	SRM3	Suc15	100
		SRM6	Fruc40	100
GS	Suc30	SRM3	Fruc30	56.67
OB	Fruc30	SRM3	Fruc40	74
FP	Suc30	SRM1	Fruc30	37.5
NB	Fruc15	SRM1	Gluc30	87.5
WP	Suc30	SRM1	Gluc15	100
		SRM3	Gluc30	100